

Ecosystems monitoring powered by environmental genomics: a review of current strategies with an implementation roadmap

Cordier, Tristan; Alonso-Saez, Laura; Apothéloz-Perret-Gentil; Aylagas, Eva; Bohan, David A.; Bouchez, Agnes; Chariton, Anthony A; Creer, Simon; Fruhe, Larissa; Keck, Francois; Keeley, Nigel; Laroche, Olivier; Leese, Florian; Pochon, Xavier; Stoeck, Thorsten; Pawlowski, Jan; Lanzen, Anders

Molecular Ecology

DOI:

<https://doi.org/10.1111/mec.15472>

Published: 01/07/2021

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Cordier, T., Alonso-Saez, L., Apothéloz-Perret-Gentil, Aylagas, E., Bohan, D. A., Bouchez, A., Chariton, A. A., Creer, S., Fruhe, L., Keck, F., Keeley, N., Laroche, O., Leese, F., Pochon, X., Stoeck, T., Pawlowski, J., & Lanzen, A. (2021). Ecosystems monitoring powered by environmental genomics: a review of current strategies with an implementation roadmap. *Molecular Ecology*, 30(13), 2937-2958. <https://doi.org/10.1111/mec.15472>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Ecosystems monitoring powered by environmental genomics: a review of current strategies with an implementation roadmap.

Tristan Cordier^{1*}, Laura Alonso-Sáez², Laure Apothéloz-Perret-Gentil¹, Eva Aylagas³, David A. Bohan⁴, Agnès Bouchez⁵, Anthony Chariton⁶, Simon Creer⁷, Larissa Frühe⁸, François Keck⁵, Nigel Keeley¹⁰, Olivier Laroche¹⁰, Florian Leese^{11,12}, Xavier Pochon^{13,14}, Thorsten Stoeck⁸, Jan Pawlowski^{1,15,16}, Anders Lanzén^{2,17}

¹Department of Genetics and Evolution, University of Geneva, Science III, 4 Boulevard d'Yvoy, 1205 Geneva, Switzerland

²AZTI, Txatxarramendi Irla s/n, 48395 Sukarrieta, Spain

³Red Sea Research Center (RSRC), Biological and Environmental Sciences and Engineering (BESE), King Abdullah University of Science and Technology (KAUST), 23955-6900 Thuwal, Saudi Arabia

⁴Agroécologie, AgroSup Dijon, INRAE, Univ. Bourgogne, Univ. Bourgogne Franche-Comté, 17 rue Sully, BP 86510, F-21000, Dijon, France

⁵UMR CARTEL, INRAE, F-74200 Thonon, France

⁶Department of Biological Sciences, Macquarie University, NSW, Australia 2113

⁷School of Natural Sciences, Bangor University, Gwynedd, LL57 2UW, UK

⁸Department of Ecology, Technische Universität Kaiserslautern, Erwin-Schroedinger-Str. 14, D-67663 Kaiserslautern, Germany

¹⁰Benthic Resources and Processes Group, Institute of Marine Research, Tromsø, Norway

¹¹Aquatic Ecosystem Research, Faculty of Biology, University of Duisburg-Essen, Universitaetsstr. 5, 45141 Essen, Germany

¹²Centre for Water and Environmental Research (ZWU), University of Duisburg-Essen, Universitaetsstr. 2, 45141 Essen, Germany

¹³Coastal & Freshwater Group, Cawthron Institute, 98 Halifax Street East, 7010 Nelson, New Zealand

¹⁴Institute of Marine Science, University of Auckland, Warkworth 0941, New Zealand

¹⁵ID-Gene ecodiagnosics, Campus Biotech Innovation Park, 1202, Geneva, Switzerland

¹⁶Institute of Oceanology, Polish Academy of Sciences, 81-712 Sopot, Poland

¹⁷IKERBASQUE, Basque Foundation for Science, Bilbao, Spain

Contact: tristan.cordier@gmail.com

Key words: Biomonitoring; metabarcoding; environmental DNA; biodiversity; implementation strategy; ecosystem management

Abstract

A decade after environmental scientists integrated high-throughput sequencing technologies in their toolbox, the genomics-based monitoring of anthropogenic impacts on the biodiversity and functioning of ecosystems is yet to be implemented by regulatory frameworks. Despite the broadly acknowledged potential of environmental genomics to this end, technical limitations and conceptual issues still stand in the way of its broad application by end-users. In addition, the multiplicity of potential implementation strategies may contribute to a perception that the routine application of this methodology is premature or “in development”, hence restraining regulators from binding these tools into legal frameworks. Here, we review recent implementations of environmental genomics-based methods, applied to the biomonitoring of ecosystems. By taking a general overview, without narrowing our perspective to particular habitats or groups of organisms, this paper aims to compare, review and discuss the strengths and limitations of four general implementation strategies of environmental genomics for monitoring: (A) Taxonomy-based analyses focused on identification of known bioindicators or described taxa; (B) *De novo* bioindicator analyses; (C) Structural community metrics including inferred ecological networks; and (D) Functional community metrics (metagenomics or metatranscriptomics). We emphasise the utility of the three latter strategies to integrate meiofauna and microorganisms that are not traditionally utilised in biomonitoring because of difficult taxonomic identification. Finally, we propose a roadmap for the implementation of environmental genomics into routine monitoring programs that leverage recent analytical advancements, while pointing out current limitations and future research needs.

The need for broad scale ecosystem monitoring strategies

Biodiversity drives the fundamental processes of ecosystems and provides invaluable services on which we depend. Anthropogenic, detrimental impacts on ecosystems, including accelerating climate change, are unprecedented (Waters et al., 2016) and have led to a decline of biodiversity across the globe (Butchart et al., 2010; Cardinale et al., 2012; Hughes et al. 2018). Recent reports stress that one out of the 8 million known species are presently at risk of extinction (IPBES report, 2019). This threatens ecosystem function(ing) and services. Therefore, the urgent challenge is now to build a set of efficient tools to enhance our capacity to predict or detect early warnings of critical ecological shifts efficiently, in order to forecast the direction of such shifts and their impacts on ecosystem functions and services (Carpenter et al., 2011; Barnosky et al., 2012; Ratajczak et al., 2018).

Because our societies aim to reach a trade-off between socioeconomic development and ecosystems sustainability (UN A/RES/70/1, 2015), regulatory frameworks have been established worldwide for the sustainable development of industries within environmental constraints (Niemeijer 2002; Grizetti et al., 2015). Such regulatory systems have been incorporated into various national and international directives, especially for aquatic ecosystems (e.g. the Water Framework Directive, WFD, Directive 2000/60/EC and Marine Strategy Framework Directive, MSFD, Directive 2008/56/EC in Europe, the Clean Water Act of the US Environmental Protection Agency in the USA, as well as the United Nations Convention on the Law of the Sea, UNCLOS). The backbone of such monitoring programs is the biological component of ecosystems, as a measure of ecosystem 'health' or 'integrity' (Karr, 1999). This biological component is often referred to as the Biological Quality Elements in those regulations (BQEs, Borja et al., 2013; Hering et al., 2018). Most monitoring strategies implemented in regulations rely on the bioindication principle (autecology, Box 1), i.e. significant correlations between the occurrence of specific organisms and a set of environmental variables. Although chemical and hydrological monitoring techniques provide an environmental quality snapshot, biological indicators convey a cumulative time-integrated measure as their occurrence is the product of their local adaptation and their responses to ecosystem variations and/or disturbances across an extended period of time (Carignan & Villard, 2002; Lear et al., 2011; Birk et al., 2012).

92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122

The limits of currently implemented ecosystem monitoring strategies

Traditionally, morphologically distinguishable invertebrates have been used as bioindicators in both aquatic and terrestrial ecosystems (Reynoldson & Metcalfe-Smith, 1992; Bongers & Ferris, 1999; Hodgkinson & Jackson, 2005; Gerlach et al., 2013). Fishes, amphibians, macrophytes, phytoplankton and diatoms, are also routinely used in aquatic ecosystems (Birk et al., 2012). Various Biotic Indices (BIs) have been formalized, based on the predictable responses of bioindicator species to environmental disturbances (autecological value) in marine (Maurer et al., 1999; Borja et al., 2000; Rygg et al., 2013), freshwater (Kelly et al., 1995; Stark et al., 1998; Prygiel & Coste, 2000) and terrestrial (Urzelai et al., 2000; Marull et al., 2007) ecosystems. Almost half of the monitoring methodologies currently used in Europe rely on such BIs (Birk et al., 2012). However, for environments or geographical regions for which no BI has been calibrated, ecological assessments rely instead on biodiversity measures of “charismatic” groups such as fishes (Pont et al., 2006), amphibians (Welsh et al., 1998) and insects (Basset et al., 2004).

Morphology-based methodologies require the collection and identification of hundreds to thousands of specimens per sample, which is a slow, labor-intensive process. These limitations seriously hamper our capacity to scale up biomonitoring and satisfy the increasing demand for environmental monitoring programs in a timely fashion that allows informed ecosystem management (Baird & Hajibabaei, 2012). Moreover, this conventional morphology-based approach is compromised by several other shortcomings: (i) it focuses only on morphologically identifiable biodiversity, ignoring the inconspicuous meiofaunal and microbial domains, which are known to include powerful bioindicators; (ii) cryptic diversity remains unrecognized (morphologically indistinguishable look-alikes with differing tolerance to disturbances); (iii) variation in species life stages, damaged specimens and misidentifications caused by decreasing taxonomic expertise worldwide may lead to variable and noisy species’ inventories, and by extension, to uncertain ecological assessments. Taken together, the need for faster, more objective, robust and cost-effective tools and strategies to deliver a more efficient ecosystem monitoring has never been more pressing.

The environmental genomics revolution for biodiversity research and ecosystem monitoring

Over the last decade, the development of environmental genomics (EG) coupled with high-throughput sequencing (HTS) technologies has led to a marked improvement in our ability to document biodiversity patterns, for both species occurrence (amplicon sequencing, i.e. metabarcoding, reviewed in Bohmann et al., 2014; Valentini et al., 2016; Deiner et al., 2017; Cristescu et al., 2018; Taberlet et al. 2018; Ruppert et al., 2019) and their metabolic functions (metagenomics and metatranscriptomics, reviewed in Ungerer et al., 2008; Vandenkoornhuysen et al., 2010; Quince et al., 2017; Singer et al., 2017; Escalas et al., 2019). Multidisciplinary teams and consortiums have initiated large scale projects aiming at collecting biodiversity data using EG throughout the globe, to address fundamental ecological questions. Among these initiatives, the large barcoding projects led by the international Barcode of Life (Ratnasingham & Hebert, 2007), the Earth Microbiome Project (Gilbert et al., 2010) and the TARA Oceans Project (Karsenti et al., 2011) represent three of the most emblematic examples. Those projects have unraveled an unexpected cryptic (Bickford et al., 2007) and novel microbial diversity (the ‘unseen majority’) guiding reconstruction of the eukaryotic tree of life (Adl et al., 2019). Even though this microbial diversity is known to represent a key component of ecosystem functioning (Delgado-Baquerizo et al., 2016; Guidi et al., 2016; Cavicchioli et al., 2019), the ecology of most microorganisms remains largely enigmatic.

The potential of EG for surveying biodiversity and monitoring natural ecosystems at a broad spatio-temporal scale was quickly identified and implemented by environmental scientists (Baird & Hajibabaei, 2012; Taberlet et al., 2012; Davies et al., 2012; Kelly et al., 2014). This work has been boosted by the massive drop in sequencing costs, with over four orders of magnitude within the last 15 years (<https://www.genome.gov>). This has enabled numerous clinical and environmental routine applications. Indeed, fueled by the continuous efforts to optimize laboratory protocols and bioinformatic tools, all steps from large-scale collection of samples, generation of HTS data, statistical analysis, and interpretation of results, can now be performed in matter of days or weeks (Juul et al., 2015; Quinn et al., 2016; Deshpande et al., 2019; Reintjes et al., 2019). For aquatic ecosystems especially, the next breakthrough of this revolution is now expected to be the development and deployment of low-cost, automated and miniaturized *in situ* environmental nucleic acids (eDNA/RNA) samplers (Carr et al., 2017; Gan et

al., 2017). These may be integrated into autonomous instruments for broad-scale and continuous ecosystem monitoring programs (Brandt et al., 2016; Bohan et al., 2017; Aguzzi et al., 2019; Benway et al., 2019; Levin et al., 2019).

These advances in genomics-based research have led to a series of pilot studies assessing the applicability of EG for the monitoring of ecosystem changes by collecting biodiversity data from various taxonomic groups (e.g. fishes, macroinvertebrates, protists, bacteria) and environments (e.g. water, biofilms, soil or sediment). Several such pilot studies have targeted multicellular organisms as a replacement for arduous morphological identification of the same taxa (Hajibabaei et al., 2011, 2012; Thomsen et al., 2012; Zhou et al., 2013; Lejzerowicz et al., 2015). However, the potential of EG to leverage the general eukaryotic and prokaryotic diversity for ecological monitoring, has also been explored (Chariton et al., 2010; Bik et al., 2012; Dowle et al., 2015; Lallias et al., 2015), and indeed advocated (Creer et al., 2010; Payne, 2013; Bouchez et al., 2016; Chariton et al., 2016; Graham et al., 2016). Encouraged by the immense opportunities for ecosystem monitoring, over 45 countries recently decided to join their efforts within the European COST Action DNAqua-Net, to anticipate upcoming paradigm-shifts and develop genomic tools tailored for the monitoring of aquatic ecosystems (<http://dnaqua.net>, Leese et al., 2016). Similarly, other large-scale collaborative projects were recently launched, including STREAM in Canada (<https://stream-dna.com/>), Lakes380 in New Zealand (<https://lakes380.com/>) and NGB in France (<http://next-genbiomonitoring.org/>), aiming at the unbridling of EG for ecosystem monitoring.

Multiple pilot and methodological EG studies have highlighted important variation in terms of compliance with current regulatory programs (reviewed in Hering et al., 2018), leading to the proposition of multiple implementation strategies for current and future ecosystem monitoring programs. Here, we compare and review the strengths and limitations of these EG-based strategies for ecosystem monitoring. Our objective is to pinpoint the criteria of existing monitoring programs that could be fulfilled by EG methods as of today, and clarify the work ahead for the monitoring programs that could benefit from EG in the near future, given continued technological and analytical advancements. To this end, we classify these strategies into four broad categories (Figure 1, Table S1): (A) Taxonomy-based analyses that focus on known bio-indicator species, or the identification and enumeration of formally or informally described taxa; (B) *De novo* bioindicator analyses aiming to identify and utilise novel

bioindicators, independent of formal taxonomy; (C) Structural community metrics relying on community structure or inferred ecological networks, where taxa are interchangeable; and (D) Functional community metrics or indicators that focus on protein-coding genes or transcripts instead of taxonomic composition. Based on the specificities of each strategy, their level of maturity and their compatibility with existing regulations (Table 1), we propose an implementation roadmap to integrate EG into ecosystems monitoring programs and highlight future research needs to be undertaken.

“Taxonomy-based” strategy: screening known species and bioindicators with environmental genomics

This strategy relies on the enumeration of known biodiversity from DNA obtained from an environmental sample (e.g. sediment, soil, biofilm, water) or from bulk material prepared from an environmental sample by e.g. elutriation, trapped individuals or biofilm scratching (Figure 1A). This strategy closely fits the conventional, morphology-based monitoring approach, because it primarily aims at reaching a satisfactory level of congruence in terms of both qualitative and quantitative biodiversity inventories. The taxonomy-based strategy is *de facto* limited to the morphologically characterized fraction of biodiversity for which reference sequences are available in public databases. Hence, approaches using it have usually overlooked meiofaunal or microbial taxa, difficult to identify on the basis of morphological traits, and for most of which the autecology is poorly known (but see Pawlowski et al., 2016). The reference databases routinely used by EG studies include for instance the universal but essentially non-curated GenBank nucleotide repository from the National Center for Biotechnology Information (Benson et al., 1999, but see Leray et al., 2019), or the curated databases BOLD for COI barcodes, primarily from animals (Ratnasingham & Hebert, 2007), SILVA for universal ribosomal markers (Quast et al., 2013), PR² for protists (Guillou et al., 2013), Diat.barcode for diatoms (Rimet et al., 2016), and Unite for fungi (Nilsson et al., 2018).

Depending on the environment assessed and the taxonomic group considered, the performance of taxonomy-based approaches varies considerably (Hering et al., 2018). Benchmarking studies comparing EG-based and conventional morphology-based taxonomic inventories (Table S1) have shown mixed degrees of congruence. For the non-invasive

detection of fish species from DNA traces in filtered marine water, the rate of success from taxonomy-based monitoring is reported near perfect (Thomsen et al., 2012; Bakker et al., 2017; but see DiBattista et al., 2017). For freshwater macroinvertebrate bulk samples, the rate of species detection varied from 67% (Elbrecht et al., 2017) to 73-83% (Hajibabaei et al., 2011; 2012). In contrast, for benthic diatoms sampled from biofilms, the congruence of morphological taxonomy and EG-inferred taxonomy, in terms of shared taxa at species level, ranged only from 15-18% (Rivera et al., 2017; Vasselon et al., 2017a) to 28% (Visco et al., 2015). The reported congruence for macroinvertebrates sampled from marine sediments ranged from 20% (Lejzerowicz et al., 2015) up to 60% (Aylagas et al., 2016). Noteworthy, those studies also detected numerous species that were unnoticed in morphological inventories (Hajibabaei et al., 2011; 2012; Elbrecht et al., 2017). Despite these discrepancies, the studies inferring BI values from the detected bioindicators species show very promising results, for both freshwater diatoms (Kermarrec et al., 2014; Visco et al., 2015; Vasselon et al., 2017b; Kelly et al., 2018) and macroinvertebrates (Elbrecht et al., 2017) as well as for marine macroinvertebrates (Lejzerowicz et al., 2015; Aylagas et al., 2016). While acknowledging that the congruence for both qualitative and quantitative inventories are not fully satisfactory, these studies have demonstrated that EG tools are still able to detect sufficient bioindicator taxa to infer accurate BI values, even when considering only presence/absence (Aylagas et al., 2016). The EG methodology has therefore been promoted as a promising tool for fast and cost-effective biodiversity screening for ecosystem monitoring, even while the simultaneous collection of classical morphological samples for validation is univocally suggested. Nonetheless, further improvements in molecular protocols as well as BI inter-calibration is a necessity towards harmonization and standardization across Europe (Poikane et al., 2014; Hering et al., 2018) and beyond (Jeunen et al., 2019).

Various biological and technical limitations still impede the implementation of the taxonomy-based strategy for routine monitoring applications (Leese et al., 2018). These limitations mainly stem from the fact that the methods sample fundamentally different units of presence (molecules *versus* individuals), resulting in different biases affecting richness, abundance and taxonomic composition. The richness of “molecular species”, i.e. Operational Taxonomic Units (OTUs) or Amplicon Sequence Variants (ASVs, the new operational unit paradigm, Callahan et al., 2017), should not be considered analogous to morpho-species

richness even in the theoretical absence of noise resulting from PCR and sequencing biases. This discrepancy is due to cryptic diversity (Stork, 2018), intragenomic or intra-specific marker variation (Bik et al., 2013, Sun et al., 2013), and the presence of DNA from dead and inactive organisms or as extracellular DNA (Collins et al., 2018). Likewise, the abundance of taxa inferred from HTS read counts can typically not be used to infer the number of individuals. Indeed, the number of sampled DNA molecules and sequence reads are a consequence of the number of individuals, but also of the biomass and the variable number of copies of the targeted marker in the genome (Bik et al., 2013, Vetrovský, et al., 2013), in addition to variations in DNA extractability and primer-specific amplification bias (Elbrecht et al., 2015; Piñol et al., 2015; Krehenwinkel et al., 2017). Finally, EG studies suffer from a strong sampling effect because DNA extractions are typically performed from small amounts of material, making large-size organisms less well-represented in eDNA extracts (Lanzén et al., 2017). However, bulk samples (Elbrecht et al., 2017), larger extraction volume (Nascimento et al., 2018) or more aggressive homogenization (Lanzén et al., unpublished data) can partially alleviate this issue.

Since the taxonomy-based strategy depends on reference sequences for organism identification, the incompleteness of reference databases can also have a major impact. Hence, completing databases, both by the “vertical” addition of more taxa and by the “horizontal” coverage of wider geographical areas, would certainly contribute to an improvement in identification (Vasselon et al., 2017; McGee et al., 2019). However, despite sustained efforts, reference databases will likely remain skewed towards some taxa, while suffering from important gaps across other taxonomic groups or biogeographical regions (Weigand et al., 2019; McGee et al., 2019). All these issues directly impact both of the key parameters for applying BIs to assess impact, namely the qualitative and quantitative measures of biodiversity (Pawlowski et al., 2018).

Nevertheless, multiple studies have shown that there is room for considerable improvements to better bridge the current gaps between taxonomy-dependent molecular and morphology-based approaches. Taxonomic breadth in HTS data could be broadened by carefully designing novel amplification primers (Elbrecht et al., 2019) or using more than one primer pair (Corse et al., 2019). Applying correction factors to read counts, based on established knowledge of the biovolume (Vasselon et al., 2018), the number of copies of the targeted marker (Vetrovský, et al., 2013) or by spiking samples with known internal standard for

quantitative determinations (Tkacz et al., 2018; Ji et al., 2019), are all promising methods for resolving these challenges. Finally, the integration of bioinformatic tools for the automated curation of databases from mislabeled sequences will improve their reliability (Ashelford et al., 2005; Kozlov et al., 2016).

“De novo” strategy: discovering new bioindicators and harnessing them for routine monitoring.

In contrast to the taxonomy-based strategy, the *de novo* one does not immediately generate an ecological assessment, because it does not employ previous knowledge associated with bioindicators. Instead, the *de novo* strategy aims at establishing new bioindicators using EG-based profiling of communities and independently generated ecological status or known disturbance gradients (Figure 1B). Harnessing EG and HTS technologies to explore a broader range of biological diversity, formally labelled or not (i.e. taxonomically described or identified), represents an opportunity to move towards a more holistic monitoring paradigm (Chariton et al., 2010; Bik et al., 2012). By considering all the OTU (or ASV) profiles along a known impact gradient of typical anthropogenic origin, studies applying this strategy have shown that HTS data represent a virtually unlimited reservoir of new bioindicators. Examples (listed in Table S1) include contamination by pesticides (Thompson et al., 2016; Andújar et al., 2017) or other agricultural stressors (Salis et al., 2017), and gradients of eutrophication and urban contamination in freshwater systems (Apothéloz-Perret-Gentil et al., 2017; Martínez-Santos et al. 2018; Simonin et al., 2019; Tapolczai et al., 2019a, 2019b). In marine environments, the utility of this strategy has been demonstrated after an oil spill (Bik et al., 2012), in the vicinity of offshore drilling platforms (Lanzén et al., 2016; Laroche et al., 2016, 2018a) and aquaculture sites (Pawlowski et al. 2014, Pochon et al., 2015; Dowle et al., 2015; Keeley et al., 2018; Stoeck et al. 2018a, 2018b) as well as along eutrophication and urban or industrial contamination gradients in estuaries (Chariton et al., 2010, 2015; Angly et al., 2015; Lallias et al., 2015; Obi et al., 2016). Interestingly, most of the studies sampling marine sediments highlighted that meiofaunal invertebrates, such as nematodes, gastrotrichs and platyhelminths (Chariton et al., 2010; Bik et al., 2012; Lanzén et al., 2016), large groups of protists such as diatoms, oomycetes and ciliates (Lanzén et al., 2016; Stoeck et al., 2018a) or foraminifera (Pawlowski et al., 2014; Laroche et al., 2016; Frontalini et al., 2018) but also fungi

(Bik et al., 2012) and bacteria (Angly et al., 2015; Dowle et al., 2015; Martínez-Santos et al. 2018; Obi et al., 2016; Aylagas et al., 2017; Stoeck et al., 2018b; Keeley et al. 2018) have great potential as bioindicators of anthropogenic impacts and can readily be captured by EG studies.

Unfortunately, most proof-of-concept studies employing the *de novo* strategy have not yet validated their results by performing ecological assessments based on newly identified bioindicators as a reference in a new environmental context. For this information to be useful on new samples, the data obtained from known disturbance gradients (i.e. reference or training dataset) must be operational in different spatiotemporal contexts. To this end, two main approaches have been proposed and tested, namely indicator value (e.g. the IndVal approach, Dufrêne and Legendre 1997) and supervised machine learning (SML, Crisci et al., 2012; Libbrecht & Noble, 2015).

The indicator value approach ascribes autecological values (or discrete “eco-groups”) to OTUs or ASVs based on their occurrence in samples of known disturbance level, in a similar manner as for the establishment of morphology-based bioindicators. Hence, the autecological values of these *de novo* bioindicators are directly calibrated on the HTS data, which alleviates the qualitative and quantitative biases encountered with the taxonomy-based EG strategy. This has proven successful for both freshwater benthic diatoms (Apothéloz-Perret-Gentil et al., 2017; Tapolczai et al., 2019a, 2019b) and for bacterial and eukaryotic communities in streams and estuarine systems (Chariton et al., 2015; Li et al., 2018). An analogous approach is the use of polynomial quantile regression splines (Andersson, 2008). This has shown great promise for the prediction of impacts from organic enrichment in aquaculture sites using eukaryotic and prokaryotic metabarcoding data in parallel (Keeley et al., 2018). For diatoms, the accuracy of the assessment can be largely improved, arguably because the indicator value approach makes use of a larger number of OTUs or ASVs, compared to an approach relying solely on their taxonomic assignments (Apothéloz-Perret-Gentil et al., 2017; Tapolczai et al. 2019a, 2019b).

Supervised machine learning (SML) also requires training datasets, i.e. reference disturbance levels (labels) associated with the community profiles of the samples (features). These algorithms are best at classification problems involving multidimensional and noisy datasets (Libbrecht & Noble, 2015), which are common attributes of HTS data. The task is to automatically disentangle the feature signal (OTU or ASV profiles) and their co-occurrence that convey an ecological signal from background noise. This extracted knowledge is self-contained

in a trained model that can be used to make predictions of disturbance level on new samples, based on their compositional profiles (Cordier et al., 2019a). Supervised machine learning also alleviates the qualitative and quantitative biases that hamper the taxonomy-based strategy in a more straightforward manner, because the model is trained directly on HTS data. The applicability of SML has been demonstrated in marine environments, for the detection of various pollutants (Smith et al., 2015) and for the prediction of aquaculture impacts on benthic biodiversity (Cordier et al., 2017; 2018). The SML-based inference of BI values has also been shown to outperform the taxonomy-based strategy, relying on the detection of established macroinvertebrates bioindicators DNA (Cordier et al., 2018), and may be more powerful than the IndVal approach (Frühe et al., 2020). Supervised machine learning applications have also succeeded in predicting the origin of container ship ballast waters (Gerhard & Gunsch, 2019).

The *de novo* strategy provides numerous advantages over the taxonomy-based one. First, it can reduce or bypass the dependence on reference sequence databases for taxonomic assignments of HTS reads to known bioindicators. Instead, new ecological knowledge is hypothesised *de novo* during the calibration of OTUs or ASVs autecological values (IndVal) or during the supervised training of a model (SML). Second, it can leverage powerful but previously inaccessible groups of bioindicators among prokaryotes, protists, meiofauna and mesozooplankton, that are widespread and may react both faster and stronger to environmental disturbances (Creer et al., 2010; Payne, 2013; Bouchez et al., 2016; Pawlowski et al. 2016). Finally, when applied for the inference of BIs that are currently employed in routine monitoring programs, a *de novo* strategy is directly compatible with current regulations, because the assessment categories remain the same and the BI values are simply inferred indirectly. Hence, this strategy assures a full backward and forward compatibility with current monitoring programs, facilitating continuity of important time series datasets (Bálint et al., 2018).

“Structural community metrics” strategy: blending theoretical ecology into routine ecosystem monitoring.

This strategy relies on metrics extracted from the community structure or from inferred ecological networks – where taxa are interchangeable – in order to assess the impact of disturbance and its ramifications on ecosystem functioning (Figure 1C). This represents a clear

paradigm-shift for ecosystem monitoring programs, because the evaluation of bioindicators, based on the compositional variation of communities, is not the main aim of the strategy. Instead, its focus is to discover and understand the ecological processes shaping biological communities and their response to disturbances, which is indeed one of the core questions of ecological research. It has long driven the exploration of the links between generic, taxonomy and composition-independent biodiversity metrics or species functional traits distribution and ecosystems functioning and resilience, to reach a more general theoretical framework (Cardinale et al., 2000; McCann, 2000; Hooper et al., 2005; Tilman et al., 2006; Ives & Carpenter, 2007; Mouillot et al., 2013; Loreau & de Mazancourt, 2013).

Structural community metrics can be computed from compositional data generated by EG studies, including alpha diversity (e.g. OTU or ASV richness, Shannon diversity or Pielou evenness; reviewed in Daly et al., 2018), along with its phylogeny-aware derivatives (reviewed in Tucker et al., 2017; Washburne et al., 2018). Under anthropogenic impact, alpha diversity in marine sediment has been found to decrease for foraminifera (Pawlowski et al., 2014; 2016; Laroche et al. 2018b), ciliates (Stoeck et al., 2018a) and bacterial communities (Stoeck et al., 2018b). Conversely, disturbances in marine sediments can also trigger increases in bacterial diversity and metabolic activity (Galand et al., 2016; Pérez-Valera et al., 2017). This suggests that the variation of alpha diversity alone is insufficient as a widely applicable indicator of disturbance. Phylogeny-aware metrics attempt to account for the evolutionary relationships among taxa composing communities, to provide insights into community assembly processes and by extension their predictable responses to environmental variations (Webb et al., 2002; Cavender-Bares et al., 2009, but see Mayfield & Levine, 2010; Gerhold et al., 2015). This relationship between phylogenetic diversity and ecosystem functioning has received a lot of attention by plant ecologists (Flynn et al., 2011). However, only few studies have employed EG data to this end, targeting mostly microbial groups, which, as for simple alpha-diversity metrics, has resulted in contrasting conclusions (Galand et al., 2015; Pérez-Valera et al., 2017, Liu et al., 2017; but see Venail & Vives, 2013; Keck & Kahlert, 2019 for studies employing sequencing data but not strictly EG).

Metrics based upon alpha diversity may be misleading (Santini et al., 2017) because their variation is often non-linear, strongly scale-dependent (Chase et al., 2019) and valuable only in comparing contexts sampled using the same methodology (Shade, 2017). It also

implicitly conveys the idea that ‘higher diversity is better’ which is not necessarily true (Shade, 2017). The inference of ecological functioning based on phylogeny-aware metrics relies on the niche conservatism concept, which postulates that closely related taxa share similar functional traits (Webb et al., 2002; Cavender-Bares et al., 2009; Srivastava et al., 2012). Under this assumption, increased phylogenetic diversity may support functionally diverse or multifunctional ecosystems (Hector & Baghi, 2007 but see Manning et al., 2018). By extension, higher phylogenetic diversity may also support ecosystem resilience, provided that the species fulfilling similar functions have differing responses to disturbances (Cadotte et al., 2012; Oliver et al., 2015). However, because not all functional traits necessarily have a phylogenetic signal (Srivastava et al., 2012), including for microbes (Martiny et al., 2013), inferring ecosystem functioning and the level of anthropogenic impact based on phylogeny-aware metrics alone may prove to be misguided. Likewise, conservation strategies based on these metrics may also be suboptimal (Mazel et al., 2018).

Another set of structural community metrics can be computed from the topology of inferred ecological or co-occurrence networks, representing potential biotic interactions (reviewed in Faust & Raes, 2012; Vacher et al., 2016; Layeghifard et al., 2017). Based on empirical evidence of the variation in network structure under environmental disturbance (Tylianakis et al., 2007; Zhou et al., 2011; Karimi et al., 2016; Ma et al., 2018), their properties have been suggested as potential indicators of ecosystem functioning and integrity (Gray et al., 2014; Karimi et al., 2017; Bohan et al., 2011, 2017; Lau et al., 2017; Tylianakis et al., 2017; Pellissier et al., 2018; Delmas et al., 2019). In recent years, a growing interest in these approaches has led to a series of studies employing EG to infer ecological networks from microbial community data (Zhou et al., 2011; Lupatini et al., 2014; Zappelini et al., 2015; Pérez-Valera et al., 2017; Pauvert et al., 2019) or from macroinvertebrates (Compson et al., 2019), in order to explore the links between network properties such as connectance, centrality or nestedness, and ecosystem functioning. For instance, it has been shown that bacterial communities in anthropized soil may have fewer potentially interacting taxa, than in natural soil (Lupatini et al., 2014). Likewise, in aquatic ecosystems, anthropogenic impacts are reflected in co-occurrence networks by a lower connectivity (Lawes et al., 2017; Laroche et al., 2018b; Li et al., 2018) and a lower ratio of positive interactions (Laroche et al., 2018b).

While promising, exploring the links between the properties of ecological networks inferred from EG data and ecosystem functioning is still in its infancy (Faust et al., 2012; 2015; Lima-Mendez et al., 2015; Lawes et al., 2017; Laroche et al., 2018b; Li et al., 2018; Pauvert et al., 2019). Multiple methodological issues limit the inference of robust networks from EG data based on co-occurrences in space or time. For example, read counts are strictly compositional, representing relative abundance of the marker itself, rather than presence or absolute abundances (but see Friedman & Alm, 2012; Kurtz et al., 2015). Further, it is challenging to control for covariates and confounding environmental parameters (but see Tammadoni-Nezhad et al., 2013; Tackmann et al., 2018; Cougoul et al., 2018; Chiquet et al., 2018; Momal et al., 2019), replicability of inference (Pauvert et al., 2019) and the relative merits of statistical and logical inference (Vacher et al., 2016). Robust networks also require considerably more replicates than are typically collected in EG studies, which increase both time and costs. Nevertheless, as more benchmark datasets containing both EG data and independently confirmed interactions between taxa become available to complement simulated datasets (see Lima-Mendez et al., 2015), making robust network inference to explore the applicability of their metrics for ecosystem monitoring will likely come within reach in the years to come.

“Functional community metrics” strategy: employing functional environmental genomics for routine monitoring.

Another avenue of implementation of EG for ecosystem monitoring is the use of shotgun metagenomics and metatranscriptomics, depicting the metabolic capabilities of the community, and the expressed genes at the moment of sampling, respectively (Figure 1D). However, ecologists have yet to disentangle the relative importance and relationship of taxonomic diversity and functional traits for ecosystems functioning (Flynn et al., 2011; Gagic et al., 2015). This is particularly true in microbial ecology with the “who’s there” *versus* “what they are doing” paradigms that often relate to the employed molecular methodologies, i.e metabarcoding *versus* metagenomics and metatranscriptomics (Xu et al., 2014). Some metagenomic contigs and functional transcripts were indeed found to represent efficient bioindicators of anthropogenic disturbances (Table S1), in terrestrial (de Menezes et al., 2012), groundwater (He et al., 2018), freshwater (Thompson et al., 2016; Cheaib et al., 2018; Falk et al., 2019) and marine environments (Kisand et al., 2012; Galand et al., 2016; Birrer et al., 2019), opening up potential

avenues for future routine ecosystem monitoring applications. Functional and taxonomic profiles may respond differently under anthropogenic disturbance (Cheaib et al., 2018), as well as under natural environmental variation (Barberà et al., 2012; Louca et al., 2016a; 2016b; Louca et al., 2018). This taxon-function decoupling paves the way towards a molecular trait-based ecology (Raes et al., 2011; Lajoie & Kembel 2019).

In an ecosystem monitoring context, functional profiles present two important features that anticipate these proxies to be more accurate than taxonomic profiles for the detection of a given environmental disturbance. First, because prokaryotes functional redundancy may be widespread (Louca et al., 2018; Pearman et al., 2019; but see Galand et al., 2018 and see Ramond et al. 2019 for protists), any given anthropogenic disturbance might trigger a similar response across multiple taxonomic groups. Under this assumption, ecosystem monitoring based on functional profiles may be less sensitive to biogeographical effects, random demographic drift, and species dispersal limitation than a monitoring strategy based on taxonomic profiles. This functional redundancy would also allow the establishment of a direct and mechanistic link between a measured functional response to a given anthropogenic disturbance. Second, because functional shifts are likely to occur prior to compositional ones, as a response of the taxa present to the disturbance, the variation of functional profiles may constitute useful early warnings for a timelier ecosystem management, especially the ones detected by means of metatranscriptomics. However, RNA molecules are reportedly less stable than genomic DNA, which would add challenging practical constraints that could preclude their implementation in routine ecosystem monitoring programs (but see Fordyce et al., 2013; Pochon et al., 2017; Cristescu, 2019; von Ammon et al. 2019). As a possible cost-effective “shortcut”, bacterial 16S rRNA profiles can be used to predict functional community profiles, based on evolutionary models (Langille et al., 2013; Aßhauer et al., 2015). Thus, 16S data could be also explored for searching potential functional bioindicators by this approach (Mukherjee et al., 2017; Laroche et al., 2018; Cordier 2020).

A roadmap for the implementation of environmental genomics for ecosystem monitoring

The emergence of standards for EG methodologies to be applied for monitoring programs.

The time lag between technological breakthroughs, the uptake by scientists and the implementation of research results into real management applications can be notoriously long. Even for clinical applications where the contributions of genomics have long been anticipated (Dulbecco, 1986; Manolio et al., 2013) and for which economic perspectives are obvious, its implementation for routine healthcare applications is considered to have started five years ago (Stark et al., 2019). This is three times faster than the average 17 years for any healthcare research (Morris et al., 2011). The emergence of consensual standards for methodological protocols and data formats for interoperable exchanges, represent the most challenging issue for the routine adoption (Stark et al., 2019).

The field of EG for ecosystems monitoring is experiencing similar issues and has yet to overcome some of the barriers to the necessary paradigm-shift in monitoring programs (Hering et al., 2018). Some of the noteworthy steps towards this goal were achieved with the widespread adoption of the MIGS, MIMARKS and MIxS standards in genomics, specifying the minimum information that should accompany any genome, marker gene sequences or any sequence (Field et al., 2008; Yilmaz et al., 2011). Now the most challenging part resides in the adoption of standardized methodologies to produce, store and analyze EG data for a given environmental setting. Given the variety of biological models and environmental matrices, reaching a consensus in the scientific community and formalizing standards appears very challenging, especially for metabarcoding (Pollock et al., 2018; Knight et al., 2018; Wilcox et al., 2018; Zinger et al., 2019) and its application to ecosystem monitoring (Cristescu & Hebert, 2018; Hering et al., 2018). Yet, these hurdles are not specific to genomics methodologies, but also exist for the morphology-based ones (Birk et al., 2012). Building robust, shared methodological standards is of course necessary and important efforts are deployed to reach this aim (Leese et al., 2018; Hering et al., 2018; Working Group CEN/TC230/WG28), for the sampling of eDNA (Dickie et al., 2018; Wilcox et al., 2018; CEN 2018a), the molecular protocols (Goldberg et al., 2016; Blackman et al., 2019) as well as for bioinformatics (Roy et al., 2018;

Knight et al., 2018), data interoperability (McDonald et al., 2012; Callahan et al., 2017) and reference databases (CEN, 2018b).

Matching the right implementation strategy to the right monitoring program.

Several monitoring programs may benefit quickly and reliably from an EG implementation, while others may require further optimization of molecular protocols or adjustments of their assessment criteria (Table 1). For instance, monitoring programs relying primarily on taxonomic inventories are still hindered by the lack of congruence between the recovered species list and their relative abundances, even though the biological and technical biases might be partially alleviated in the future. Furthermore, despite the sustained effort, reference sequence databases for barcoding remain skewed toward some groups and geographical locations (Weigand et al., 2019; McGee et al., 2019), limiting congruence between EG and morpho-taxonomic inventories. Hence, the taxonomy-based implementation strategy for these monitoring programs will require improvements of molecular protocols and reference databases, to generate EG data that better fit the current standards, or an adaptation of the currently implemented assessment criteria to fit the specificities of EG data (Hering et al., 2018).

Monitoring programs relying on the screening of established bioindicators for the computation of BI values are proposed as being compatible with an implementation of EG (Hering et al., 2018; Pawlowski et al., 2018). Indeed, this compatibility is greatly facilitated by the fact that the assessment criteria, i.e. BIs, are not meant to strictly rely on taxonomic inventories but rather on the autecology of bioindicators. Hence, for the taxonomy-based strategy, the BI formulations can compensate the impact of taxonomic mismatches between morphology and EG and databases incompleteness to some extent, because multiple taxa are ascribed identical autecological values, conveying similar ecological signal (Keck et al., 2018). The applicability of this approach has been demonstrated in freshwater (Elbrecht et al., 2017; Vasselon et al., 2017b; Kelly et al., 2018; Mortagua et al., 2019; Rivera et al., 2020) and in marine environments (Lejzerowicz et al., 2015; Aylagas et al., 2016). However, those studies have also shown that a large amount of sequences are not taxonomically assigned and currently omitted for ecological assessment, opening the door to new approaches that could extract ecological information from those unlabeled sequences.

The *de novo* strategy uses the occurrence of previously scrutinized sequences in samples of known BI values or other impact measures to ascribe autecological values to sequences directly, or generate a predictive model (Apothéloz-Perret-Gentil et al., 2017; Cordier et al., 2017; Tapolczai et al. 2019). Hence, these approaches are less sensitive to the biological and technical issues mentioned above, because the ecological signal (autecology) is calibrated directly on the specificities of EG data. From an implementation perspective, this *de novo* strategy thus may represent the most direct path towards implementation of EG into monitoring programs relying on BIs (Figure 2). Though somewhat unintuitive, this is because inferred BI values with a *de novo* strategy convey the same ecological meaning as they do with current methodologies, which is not the case when BIs values are inferred from bioindicators composition profiles depicted by EG data, as their autecological values were calibrated only on morphology-based data. Thus, the *de novo* strategy assures a better continuity with previous BIs data and time series and expand the range of possible bioindicators to virtually any taxa or sequence.

Structural and functional community metrics represent alternative implementation strategies that may ultimately lead to a more generic, broadly applicable ecological monitoring framework (Bohan et al., 2017; Karimi et al., 2017; Tylanakis et al., 2017; Quince et al., 2017; Singer et al., 2017; Pellissier et al., 2018; Escalas et al., 2019). These strategies hold the potential to provide a more mechanistic and functional understanding of the response of biological communities to ecosystem variation. Such knowledge could hence be included in predictive models to forecast shifts in biodiversity structure and possibly their consequences on their associated ecosystem services under different disturbance scenarios. However, an operational ecosystem monitoring framework remains to be built upon this theoretical ecological work (Figure 2), that has only partially been experimentally validated (but see Laroche et al., 2018; Ma et al., 2019). In addition, the extraction of structural or functional community metrics remain active fields of ecological research, and the emergence of a molecular trait-based ecology using metagenomics and metatranscriptomics profiles is in its infancy (Lajoie et al., 2019). Hence, it is premature to discuss their operational implementation and regulatory establishment, but their ecological benefit should be anticipated. Nevertheless, the collected labelled datasets including samples for the production of EG data in the course of future ecosystem monitoring campaigns will certainly contribute to move these possibilities forward.

Collecting reference data and eDNA/eRNA samples in parallel.

If EG-based methods are to complement or replace current morphology-based ones, the prerequisite is to establish whether they can provide similar ecological diagnostics, to ensure a smooth implementation and compatibility with existing time series (Leese et al., 2016; Bálint et al., 2018). This inevitably implies extensive parallel sampling of currently implemented and EG methodologies for some time, to build reference datasets on which the applicability can be assessed and the calibration with previous methodology performed (Leese et al., 2016; Keeley et al. 2018). To be reliable, such reference datasets have to cover a broad range of possible environmental conditions for a given ecosystem across multiple spatiotemporal scales, ideally in a balanced manner, to account for biotic interactions, random demographic drift and dispersal limitations that may interact with the anthropogenic pressures in the assembly of communities.

The collection of reference data raises concerns regarding the substantial financial investment necessary for monitoring programs adopting one or a combination of EG strategies, *versus* the “risk” of technological novelty and/or paradigm-shift. However, the collected reference datasets would still be extremely valuable in such case, because the extracted DNA/RNA alongside the accompanying reference metadata can be safely stored and re-analysed later on, assuring a forward compatibility to the limit of availability of stored DNA/RNA material (Hering et al., 2018; Jarman et al., 2018). Indeed, molecular costs are usually far less prohibitive than those related to field sampling and metadata collection. Hence, such fully labelled datasets will constitute the ideal benchmarks against which to assess the validity of any new implementation strategy based on novel technology or new paradigm.

Conclusion and further research needs

The potential for EG-based methods for ecosystems monitoring is enormous and can presently fulfil most of the requirements of current monitoring programs. Moving towards a routine use of EG is certainly a paradigm-shift, but this technological breakthrough will

overcome the limitations of current morpho-taxonomy methodologies and enable the required up-scaling to meet monitoring needs in a changing world. Without doubts, EG-based methods will pave the way for a more cost-effective, faster, reproducible and semi-automatable ecosystem monitoring framework. Regardless of the implementation strategy envisioned, the following key technological, scientific and societal improvements will be beneficial for a smoother transition:

- A collaborative and transdisciplinary design of monitoring campaigns, involving both experts, stakeholders and regulators would allow monitoring programs to more easily bridge the science-policy gap.
- A collection of reference morphological and molecular data in parallel, at least in a subset of reference points or during a transition period, will assure backward and forward compatibility of time series datasets, regardless of the envisioned implementation strategy to be decided in future monitoring campaigns.
- The efforts to complete reference sequence databases need to be sustained, by adding more representatives of the known biodiversity, with a wider geographical coverage.
- A reference database framework for *de novo* strategies needs to be established. A key requirement is the ability to reliably compare OTUs or ASVs identified in monitoring programs to formally establish knowledge about their sensitivity to disturbance.
- The taxonomic resolution level (haplotype, species, genus, family, order, class) at which HTS reads are most informative as genetic bioindicators for a given situation remains to be identified.
- For the identification of novel genetic bioindicators in complex communities, it will be important to distinguish the effect of natural (seasonal) variation from disturbance-induced community changes with rigorous experimental designs.
- Basic and replicable research is highly needed to develop a structural and functional community metrics-based implementation strategy. Such effort will likely contribute to the establishment of a more broadly applicable monitoring framework and less constrained by the database and geographical coverage limitations.

Box 1: Glossary of terms used in this paper

- Implementation strategy: Refers to the way environmental genomics data is produced and analysed in an ecosystem monitoring context. It includes the choice of all the molecular biology steps, i.e. targeted molecules (DNA versus RNA), metabarcoding (amplicon sequencing) versus metagenomics or metatranscriptomics (shotgun sequencing), and the computational biology steps (analytical approach), i.e. focusing on the taxonomically assigned sequences or considering all the sequences, the use of compositional turnovers (beta-diversity), structural metrics (alpha or phylogenetic diversity and ecological network properties) or functional metrics (functional genes or transcripts diversity).
- Environmental genomics: Suite of molecular tools to sample, process and analyse nucleic acids from an environmental sample (soil, water, sediment, feces)
- Environmental DNA/RNA: Nucleic acids present in an environmental sample. It encompasses the DNA/RNA within living multi or unicellular organisms, dead or decaying as well as extracellular material.
- Metabarcoding: A molecular workflow to simultaneously study the diversity of PCR-selected organisms from environmental samples using high-throughput sequencing. This is equivalent to amplicon sequencing of a taxonomic marker.
- Metagenomics: Shotgun sequencing of the genomic DNA isolated from an environmental sample. There is no PCR selection of particular taxonomic group and include coding as well as non-coding genomic material.
- Metatranscriptomics: Shotgun sequencing of retro-transcribed RNA isolated from an environmental sample. As for metagenomics, there is no PCR selection but includes only transcribed RNA (mRNA, rRNA), supposedly functional.
- Bioindicator: A taxon, marker sequence, gene or transcript that is used as an indicator of the ecological status of an environment.
- Autecological value: Ecological knowledge about the distribution and abundance of particular species obtained by studying interactions of individual organisms with their environments.
- Biotic Indices: Continuous or discrete variables that measure the level of disturbance of an environment based on the composition and relative abundance of bioindicator taxa (or OTUs/ASVs). Around half of the existing monitoring programs rely on biotic indices (BIs). The BIs usually includes several ordered discrete classes, usually from 'poor' to 'high' ecological status.

- Ecological network: Representation of statistically inferred biotic interactions through spatial or temporal co-occurrence or co-exclusion. Taxa (nodes) are connected by pairwise links (edges). Network ecology aims to understand how these network properties are linked to the functioning of ecosystems.

Figures and tables

Figure 1: Overview of the current methodology for the monitoring of ecosystems, that relies mostly on the morphological identification of biodiversity and / or bioindicators of anthropogenic impacts. Ecological diagnostics are performed based on reference biodiversity or on reference biotic indices for a given ecosystem. The development of environmental genomics methodologies has led to the proposition of multiple implementation strategies that can intervene at different levels of the monitoring workflow, to produce an ecological diagnostic. Green colors and smileys within boxes indicate reference or “high” ecological status while red colors and smileys represent non-reference biodiversity or “poor” ecological status (i.e. impacted environments). The colors on tags besides organisms or sequences indicate their bio-indication value (red: indicator of impact, yellow: indicator of intermediate status, green: indicator of good status). In this review paper, these strategies have been grouped in four broad categories: (A) Taxonomy-based analyses focused on identification of known bio-indicators or described taxa; (B) *De novo* bioindicator analyses; (C) Structural community metrics including inferred ecological networks; and (D) Functional community metrics (metagenomics or metatranscriptomics).

Figure 2: Strengths and limitations of the currently envisioned implementation strategies of environmental genomics for the monitoring of ecosystems, and their ability to fulfill the criteria of existing monitoring programs.

Table 1: Comparison of the four implementation strategies in terms of compatibility with current standards, backward and forward compatibility, performance, biodiversity coverage, generalization potential and ease of standardization

Table S1: List of studies employing environmental genomics for ecosystem monitoring sorted by strategy, ecosystem, targeted taxonomic group and objective.

Acknowledgements

TC, JP and LAPG were supported by the Swiss National Science Foundation (grant 31003A_179125). TC, JP, LAPG and AB were supported by the European Cross-Border Cooperation Program (Interreg France-Switzerland 2014-2020, SYNAQUA project). LAS was funded by a 'Ramón y Cajal' contract (RYC-2012-11404) from the Spanish Ministry of Economy and Competitiveness. EA is funded by the Saudi Aramco-KAUST Center for Marine Environmental Observations. DAB would like to acknowledge the financial support of the French Agence Nationale de la Recherche project NGB (ANR-17-CE32-011) and the ERA-NET C-IPM BioAWARE. AB and FK were supported by the Office Français de la Biodiversité (OFB). SC benefitted from the UK Natural Environment Research Council Grants NE/N003756/1 and NE/N006216/1. TS and LF were supported by the German Science Foundation (DFG) under grant STO414/15-1. XP is supported by the New Zealand Ministry for Business, Innovation and Employment contracts CAWX1904 (Biosecurity Toolbox) and C05X1707 (Lakes380). AB and FL were supported by DNAqua-Net COST Action CA15219 'Developing new genetic tools for bioassessment of aquatic ecosystems in Europe' funded by the European Union. AL is supported by IKERBASQUE (Basque Foundation for Science) and the Basque Government (project microgAMBI). We thank the members of DNAqua-net COST Action for helpful discussions.

References

- Adl, S. M., Bass, D., Lane, C. E., Lukeš, J., Schoch, C. L., Smirnov, A., ... Zhang, Q. (2019). Revisions to the classification, nomenclature, and diversity of Eukaryotes. *Journal of Eukaryotic Microbiology*, 66(1), 4–119. doi:10.1111/jeu.12691
- Aguzzi, J., Chatzievangelou, D., Marini, S., Fanelli, E., Danovaro, R., Flögel, S., ... Company, J. B. (2019). New high-tech flexible networks for the monitoring of deep-sea ecosystems. *Environmental Science and Technology*, 53(12), 6616–6631. doi:10.1021/acs.est.9b00409
- Anderson, M. J. (2008). Animal-sediment relationships re-visited: Characterising species' distributions along an environmental gradient using canonical analysis and quantile regression splines. *Journal of Experimental Marine Biology and Ecology*, 366(1–2), 16–27. doi:10.1016/j.jembe.2008.07.006
- Andújar, C., Arribas, P., Gray, C., Bruce, C., Woodward, G., Yu, D. W., & Vogler, A. P. (2018). Metabarcoding of freshwater invertebrates to detect the effects of a pesticide spill. *Molecular Ecology*, 27(1), 146–166. doi:10.1111/mec.14410
- Angly, F. E., Heath, C., Morgan, T. C., Tonin, H., Rich, V., Schaffelke, B., ... Tyson, G. W. (2016). Marine microbial communities of the Great Barrier Reef lagoon are influenced by riverine floodwaters and seasonal weather events. *PeerJ*, 2016(4), e1511. doi:10.7717/peerj.1511
- Apothéloz-Perret-Gentil, L., Cordonier, A., Straub, F., Iseli, J., Esling, P., & Pawlowski, J. (2017). Taxonomy-free molecular diatom index for high-throughput eDNA biomonitoring. *Molecular Ecology Resources*, 17(6), 1231–1242. doi:10.1111/1755-0998.12668
- Ashelford, K. E., Chuzhanova, N. A., Fry, J. C., Jones, A. J., & Weightman, A. J. (2005). At least 1 in 20 16S rRNA sequence records currently held in public repositories is estimated to contain substantial anomalies. *Applied and Environmental Microbiology*, 71(12), 7724–7736. doi:10.1128/AEM.71.12.7724-7736.2005
- Aßhauer, K. P., Wemheuer, B., Daniel, R., & Meinicke, P. (2015). Tax4Fun: Predicting functional profiles from metagenomic 16S rRNA data. *Bioinformatics*, 31(17), 2882–2884. doi:10.1093/bioinformatics/btv287

774 Aylagas, E., Borja, Á., Irigoien, X., & Rodríguez-Ezpeleta, N. (2016). Benchmarking DNA
 775 metabarcoding for biodiversity-based monitoring and assessment. *Frontiers in Marine*
 776 *Science*, 3(96). doi:10.3389/fmars.2016.00096
 777 Aylagas, E., Borja, Á., Muxika, I., & Rodríguez-Ezpeleta, N. (2018). Adapting metabarcoding-
 778 based benthic biomonitoring into routine marine ecological status assessment networks.
 779 *Ecological Indicators*, 95, 194–202. doi:10.1016/j.ecolind.2018.07.044
 780 Aylagas, E., Borja, Á., Tangherlini, M., Dell’Anno, A., Corinaldesi, C., Michell, C. T., ... Rodríguez-
 781 Ezpeleta, N. (2017). A bacterial community-based index to assess the ecological status of
 782 estuarine and coastal environments. *Marine Pollution Bulletin*, 114(2), 679–688.
 783 doi:10.1016/j.marpolbul.2016.10.050
 784 Bagley, M., Pilgrim, E., Knapp, M., Yoder, C., Santo Domingo, J., & Banerji, A. (2019). High-
 785 throughput environmental DNA analysis informs a biological assessment of an urban
 786 stream. *Ecological Indicators*, 104, 378–389. doi:10.1016/j.ecolind.2019.04.088
 787 Baird, D. J., & Hajibabaei, M. (2012). Biomonitoring 2.0: A new paradigm in ecosystem
 788 assessment made possible by next-generation DNA sequencing. *Molecular Ecology*, 21(8),
 789 2039–2044. doi:10.1111/j.1365-294X.2012.05519.x
 790 Bakker, J., Wangenstein, O. S., Chapman, D. D., Boussarie, G., Buddo, D., Guttridge, T. L., ...
 791 Mariani, S. (2017). Environmental DNA reveals tropical shark diversity in contrasting levels
 792 of anthropogenic impact. *Scientific Reports*, 7(1), 16886. doi:10.1038/s41598-017-17150-2
 793 Baldwin, D. S., Colloff, M. J., Rees, G. N., Chariton, A. A., Watson, G. O., Court, L. N., ... Hardy, C.
 794 M. (2013). Impacts of inundation and drought on eukaryote biodiversity in semi-arid
 795 floodplain soils. *Molecular Ecology*, 22(6), 1746–1758. doi:10.1111/mec.12190
 796 Bálint, M., Pfenninger, M., Grossart, H. P., Taberlet, P., Vellend, M., Leibold, M. A., ... Bowler, D.
 797 (2018). Environmental DNA time series in ecology. *Trends in Ecology and Evolution*, 33(12),
 798 945–957. doi:10.1016/j.tree.2018.09.003
 799 Banerji, A., Bagley, M., Elk, M., Pilgrim, E., Martinson, J., & Santo Domingo, J. (2018). Spatial and
 800 temporal dynamics of a freshwater eukaryotic plankton community revealed via 18S rRNA
 801 gene metabarcoding. *Hydrobiologia*, 818(1), 71–86. doi:10.1007/s10750-018-3593-0
 802 Barberán, A., Fernández-Guerra, A., Bohannan, B. J. M., & Casamayor, E. O. (2012). Exploration

803 of community traits as ecological markers in microbial metagenomes. *Molecular Ecology*,
804 21(8), 1909–1917. doi:10.1111/j.1365-294X.2011.05383.x

805 Barnosky, A. D., Hadly, E. A., Bascompte, J., Berlow, E. L., Brown, J. H., Fortelius, M., ... Smith, A.
806 B. (2012). Approaching a state shift in Earth's biosphere. *Nature*, 486(7401), 52–58.
807 doi:10.1038/nature11018

808 Basset, Y., Mavoungou, J. F., Mikissa, J. B., Missa, O., Miller, S. E., Kitching, R. L., & Alonso, A.
809 (2004). Discriminatory power of different arthropod data sets for the biological monitoring
810 of anthropogenic disturbance in tropical forests. *Biodiversity and Conservation*, 13(4), 709–
811 732.

812 Beazley, M. J., Martinez, R. J., Rajan, S., Powell, J., Piceno, Y. M., Tom, L. M., ... Sobecky, P. A.
813 (2012). Microbial community analysis of a coastal salt marsh affected by the Deepwater
814 Horizon oil spill. *PLoS ONE*, 7(7), e41305. doi:10.1371/journal.pone.0041305

815 Beng, K. C., Tomlinson, K. W., Shen, X. H., Surget-Groba, Y., Hughes, A. C., Corlett, R. T., & Slik, J.
816 W. F. (2016). The utility of DNA metabarcoding for studying the response of arthropod
817 diversity and composition to land-use change in the tropics. *Scientific Reports*, 6, 24965.
818 doi:10.1038/srep24965

819 Benson, D. A., Boguski, M. S., Lipman, D. J., Ostell, J., Ouellette, B. F. F., Rapp, B. A., & Wheeler,
820 D. L. (1999). GenBank. *Nucleic Acids Research*, 27(1), 12–17. doi:10.1093/nar/27.1.12

821 Benway, H. M., Lorenzoni, L., White, A. E., Fiedler, B., Levine, N. M., Nicholson, D. P., ... Letelier,
822 R. M. (2019). Ocean time series observations of changing marine ecosystems: An era of
823 integration, synthesis, and societal applications. *Frontiers in Marine Science*, 6(393).
824 doi:10.3389/fmars.2019.00393

825 Bickford, D., Lohman, D. J., Sodhi, N. S., Ng, P. K. L., Meier, R., Winker, K., ... Das, I. (2007).
826 Cryptic species as a window on diversity and conservation. *Trends in Ecology and*
827 *Evolution*, 22(3), 148–155. doi:10.1016/j.tree.2006.11.004

828 Bik, H. M., Fournier, D., Sung, W., Bergeron, R. D., & Thomas, W. K. (2013). Intra-genomic
829 variation in the ribosomal repeats of nematodes. *PLoS ONE*, 8(10), e78230.
830 doi:10.1371/journal.pone.0078230

831 Bik, H. M., Halanaych, K. M., Sharma, J., & Thomas, W. K. (2012). Dramatic shifts in benthic

832 microbial eukaryote communities following the deepwater horizon oil spill. *PLoS ONE*, 7(6),
833 e38550. doi:10.1371/journal.pone.0038550

834 Birk, S., Bonne, W., Borja, A., Brucet, S., Courrat, A., Poikane, S., ... Hering, D. (2012). Three
835 hundred ways to assess Europe's surface waters: An almost complete overview of
836 biological methods to implement the Water Framework Directive. *Ecological Indicators*,
837 18, 31–41. doi:10.1016/j.ecolind.2011.10.009

838 Birrer, S. C., Dafforn, K. A., Sun, M. Y., Williams, R. B. H., Potts, J., Scanes, P., ... Johnston, E. L.
839 (2019). Using meta-omics of contaminated sediments to monitor changes in pathways
840 relevant to climate regulation. *Environmental Microbiology*, 21(1), 389–401.
841 doi:10.1111/1462-2920.14470

842 Blackman, R., Mächler, E., Altermatt, F., Arnold, A., Beja, P., Boets, P., ... Deiner, K. (2019).
843 Advancing the use of molecular methods for routine freshwater macroinvertebrate
844 biomonitoring – the need for calibration experiments. *Metabarcoding and Metagenomics*,
845 3, 49–57. doi:10.3897/mbmg.3.34735

846 Bohan, D. A., Caron-Lormier, G., Muggleton, S., Raybould, A., & Tamaddoni-Nezhad, A. (2011).
847 Automated discovery of food webs from ecological data using logic-based machine
848 learning. *PLoS ONE*, 6(12), e29028. doi:10.1371/journal.pone.0029028

849 Bohan, D. A., Vacher, C., Tamaddoni-Nezhad, A., Raybould, A., Dumbrell, A. J., & Woodward, G.
850 (2017). Next-generation global biomonitoring: large-scale, automated reconstruction of
851 ecological networks. *Trends in Ecology and Evolution*, 32(7), 477–487.
852 doi:10.1016/j.tree.2017.03.001

853 Bohmann, K., Evans, A., Gilbert, M. T. P., Carvalho, G. R., Creer, S., Knapp, M., ... de Bruyn, M.
854 (2014). Environmental DNA for wildlife biology and biodiversity monitoring. *Trends in*
855 *Ecology and Evolution*, 29(6), 358–367. doi:10.1016/j.tree.2014.04.003

856 Bongers, T., & Ferris, H. (1999). Nematode community structure as a bioindicator in
857 environmental monitoring. *Trends in Ecology and Evolution*, 14(6), 224–228.
858 doi:10.1016/S0169-5347(98)01583-3

859 Borja, A., Franco, J., & Pérez, V. (2000). A marine biotic index to establish the ecological quality
860 of soft-bottom benthos within European estuarine and coastal environments. *Marine*

861 *Pollution Bulletin*, 40(12), 1100–1114. doi:10.1016/S0025-326X(00)00061-8

862 Borja, Angel, Elliott, M., Andersen, J. H., Cardoso, A. C., Carstensen, J., Ferreira, J. G., ...

863 Zampoukas, N. (2013). Good environmental status of marine ecosystems: What is it and

864 how do we know when we have attained it? *Marine Pollution Bulletin*, 76(1–2), 16–27.

865 doi:10.1016/j.marpolbul.2013.08.042

866 Borrell, Y. J., Miralles, L., Do Huu, H., Mohammed-Geba, K., & Garcia-Vazquez, E. (2017). DNA in

867 a bottle - Rapid metabarcoding survey for early alerts of invasive species in ports. *PLoS*

868 *ONE*, 12(9), e0183347. doi:10.1371/journal.pone.0183347

869 Bouchez, T., Blieux, A. L., Dequiedt, S., Domaizon, I., Dufresne, A., Ferreira, S., ... Ranjard, L.

870 (2016). Molecular microbiology methods for environmental diagnosis. *Environmental*

871 *Chemistry Letters*, 14(4), 423–441. doi:10.1007/s10311-016-0581-3

872 Brandt, A., Gutt, J., Hildebrandt, M., Pawlowski, J., Schwendner, J., Soltwedel, T., & Thomsen, L.

873 (2016). Cutting the umbilical: New technological perspectives in benthic deep-sea

874 research. *Journal of Marine Science and Engineering*, 4(2), 36. doi:10.3390/jmse4020036

875 Bucklin, A., Yeh, H. D., Questel, J. M., Richardson, D. E., Reese, B., Copley, N. J., & Wiebe, P. H.

876 (2019). Time-series metabarcoding analysis of zooplankton diversity of the NW Atlantic

877 continental shelf. *ICES Journal of Marine Science*, 76(4), 1162–1176.

878 doi:10.1093/icesjms/fsz021

879 Butchart, S. H. M., Walpole, M., Collen, B., Van Strien, A., Scharlemann, J. P. W., Almond, R. E.

880 A., ... Watson, R. (2010). Global biodiversity: Indicators of recent declines. *Science*,

881 328(5982), 1164–1168. doi:10.1126/science.1187512

882 Cadotte, M. W., Dinnage, R., & Tilman, D. (2012). Phylogenetic diversity promotes ecosystem

883 stability. *Ecology*, 93(8 SPEC. ISSUE), 223–233.

884 Callahan, B. J., McMurdie, P. J., & Holmes, S. P. (2017). Exact sequence variants should replace

885 operational taxonomic units in marker-gene data analysis. *ISME Journal*, 11(12), 2639–

886 2643. doi:10.1038/ismej.2017.119

887 Cardinale, B. J., Nelson, K., & Palmer, M. A. (2000). Linking species diversity to the functioning of

888 ecosystems: On the importance of environmental context. *Oikos*, 91(1), 175–183.

889 doi:10.1034/j.1600-0706.2000.910117.x

890 Cardinale, Bradley J., Duffy, J. E., Gonzalez, A., Hooper, D. U., Perrings, C., Venail, P., ... Naeem,
891 S. (2012). Biodiversity loss and its impact on humanity. *Nature*, 486(7401), 59–67.
892 doi:10.1038/nature11148

893 Carignan, V., & Villard, M. A. (2002). Selecting indicator species to monitor ecological integrity:
894 A review. *Environmental Monitoring and Assessment*, 78(1), 45–61.
895 doi:10.1023/A:1016136723584

896 Carpenter, S. R., Cole, J. J., Pace, M. L., Batt, R., Brock, W. A., Cline, T., ... Weidel, B. (2011). Early
897 warnings of regime shifts: A whole-ecosystem experiment. *Science*, 332(6033), 1079–1082.
898 doi:10.1126/science.1203672

899 Carr, C. E., Mojarro, A., Hachey, J., Saboda, K., Tani, J., Bhattaru, S. A., ... Ruvkun, G. (2017).
900 Towards in situ sequencing for life detection. *IEEE Aerospace Conference Proceedings*, 1–
901 18. doi:10.1109/AERO.2017.7943896

902 Cavender-Bares, J., Kozak, K. H., Fine, P. V. A., & Kembel, S. W. (2009). The merging of
903 community ecology and phylogenetic biology. *Ecology Letters*, 12(7), 693–715.
904 doi:10.1111/j.1461-0248.2009.01314.x

905 Cavicchioli, R., Ripple, W. J., Timmis, K. N., Azam, F., Bakken, L. R., Baylis, M., ... Webster, N. S.
906 (2019). Scientists' warning to humanity: microorganisms and climate change. *Nature*
907 *Reviews Microbiology*, 17, 569–586. doi:10.1038/s41579-019-0222-5

908 CEN. (2018a). *Water quality - Technical report for the management of diatom barcodes*.
909 *Technical Report TR17244*.

910 CEN. (2018b). *Water quality - Technical report for the routine sampling of benthic diatoms from*
911 *rivers and lakes adapted for metabarcoding analyses. Technical Report TR17245*.

912 Chariton, A. A., Sun, M., Gibson, J., Webb, J. A., Leung, K. M. Y., Hickey, C. W., & Hose, G. C.
913 (2016). Emergent technologies and analytical approaches for understanding the effects of
914 multiple stressors in aquatic environments. *Marine and Freshwater Research*, 67(4), 414–
915 428. doi:10.1071/MF15190

916 Chariton, Anthony A., Court, L. N., Hartley, D. M., Colloff, M. J., & Hardy, C. M. (2010). Ecological
917 assessment of estuarine sediments by pyrosequencing eukaryotic ribosomal DNA.
918 *Frontiers in Ecology and the Environment*, 8(5), 233–238. doi:10.1890/090115

919 Chariton, Anthony A., Ho, K. T., Proestou, D., Bik, H., Simpson, S. L., Portis, L. M., ... Matthews,
 920 R. A. (2014). A molecular-based approach for examining responses of eukaryotes in
 921 microcosms to contaminant-spiked estuarine sediments. *Environmental Toxicology and*
 922 *Chemistry*, 33(2), 359–369. doi:10.1002/etc.2450
 923 Chariton, Anthony A., Stephenson, S., Morgan, M. J., Steven, A. D. L., Colloff, M. J., Court, L. N.,
 924 & Hardy, C. M. (2015). Metabarcoding of benthic eukaryote communities predicts the
 925 ecological condition of estuaries. *Environmental Pollution*, 203, 165–174.
 926 doi:10.1016/j.envpol.2015.03.047
 927 Chase, J. M., McGill, B. J., Thompson, P. L., Antão, L. H., Bates, A. E., Blowes, S. A., ... O'Connor,
 928 M. (2019). Species richness change across spatial scales. *Oikos*, 128(8), 1079–1091.
 929 doi:10.1111/oik.05968
 930 Cheaib, B., Boulch, M. Le, Mercier, P. L., & Derome, N. (2018). Taxon-function decoupling as an
 931 adaptive signature of Lake Microbial metacommunities under a chronic polymetallic
 932 pollution gradient. *Frontiers in Microbiology*, 9(MAY), 1–17. doi:10.3389/fmicb.2018.00869
 933 Chiquet, J., Mariadassou, M., & Robin, S. (2018). Variational inference for sparse network
 934 reconstruction from count data. In *Proceedings of the 36th International Conference on*
 935 *Machine Learning* (p. PMLR 97:1162-1171). Retrieved from
 936 <http://arxiv.org/abs/1806.03120>
 937 Coelho, F. J. R. C., Cleary, D. F. R., Costa, R., Ferreira, M., Polónia, A. R. M., Silva, A. M. S., ...
 938 Gomes, N. C. M. (2016). Multitaxon activity profiling reveals differential microbial response
 939 to reduced seawater pH and oil pollution. *Molecular Ecology*, 25(18), 4645–4659.
 940 doi:10.1111/mec.13779
 941 Collins, R. A., Wangenstein, O. S., O’Gorman, E. J., Mariani, S., Sims, D. W., & Genner, M. J.
 942 (2018). Persistence of environmental DNA in marine systems. *Communications Biology*,
 943 1(1), 1–11. doi:10.1038/s42003-018-0192-6
 944 Compson, Z. G., Monk, W. A., Hayden, B., Bush, A., O’Malley, Z., Hajibabaei, M., ... Baird, D. J.
 945 (2019). Network-based biomonitoring: exploring freshwater food webs with stable isotope
 946 analysis and DNA metabarcoding. *Frontiers in Ecology and Evolution*, 7(395).
 947 doi:10.3389/FEVO.2019.00395

948 Cordier, T. (2020). Bacterial communities' taxonomic and functional turnovers both accurately
 949 predict marine benthic ecological quality status. *Environmental DNA*, 2, 175–183.
 950 doi:10.1002/edn3.55

951 Cordier, T., Esling, P., Lejzerowicz, F., Visco, J., Ouadahi, A., Martins, C., ... Pawlowski, J. (2017).
 952 Predicting the ecological quality status of marine environments from eDNA metabarcoding
 953 data using supervised machine learning. *Environmental Science and Technology*, 51(16),
 954 9118–9126. doi:10.1021/acs.est.7b01518

955 Cordier, T., Forster, D., Dufresne, Y., Martins, C. I. M., Stoeck, T., & Pawlowski, J. (2018).
 956 Supervised machine learning outperforms taxonomy-based environmental DNA
 957 metabarcoding applied to biomonitoring. *Molecular Ecology Resources*, 18(6), 1381–1391.
 958 doi:10.1111/1755-0998.12926

959 Cordier, T., Frontalini, F., Cermakova, K., Apothéloz-Perret-Gentil, L., Treglia, M., Scantamburlo,
 960 E., ... Pawlowski, J. (2019). Multi-marker eDNA metabarcoding survey to assess the
 961 environmental impact of three offshore gas platforms in the North Adriatic Sea (Italy).
 962 *Marine Environmental Research*, 146(April 2019), 24–34.
 963 doi:10.1016/j.marenvres.2018.12.009

964 Cordier, T., Lanzén, A., Apothéloz-Perret-Gentil, L., Stoeck, T., & Pawlowski, J. (2019). Embracing
 965 environmental genomics and machine learning for routine biomonitoring. *Trends in*
 966 *Microbiology*, 27(5), 387–397. doi:10.1016/j.tim.2018.10.012

967 Corse, E., Tougard, C., Archambaud-Suard, G., Agnèse, J. F., Messu Mandeng, F. D., Bilong
 968 Bilong, C. F., ... Dubut, V. (2019). One-locus-several-primers: A strategy to improve the
 969 taxonomic and haplotypic coverage in diet metabarcoding studies. *Ecology and Evolution*,
 970 9(8), 4603–4620. doi:10.1002/ece3.5063

971 Cougoul, A. P., Bailly, X., & Wit, E. C. (2019). MAGMA: inference of sparse microbial association
 972 networks. *BioRxiv*, 538579. doi:10.1101/538579

973 Creer, S., Fonseca, V. G., Porazinska, D. L., Giblin-Davis, R. M., Sung, W., Power, D. M., ...
 974 Thomas, W. K. (2010). Ultrasequencing of the meiofaunal biosphere: Practice, pitfalls and
 975 promises. *Molecular Ecology*, 19(SUPPL. 1), 4–20. doi:10.1111/j.1365-294X.2009.04473.x

976 Crisci, C., Ghattas, B., & Perera, G. (2012). A review of supervised machine learning algorithms

977 and their applications to ecological data. *Ecological Modelling*, 240, 113–122.
 978 doi:<http://dx.doi.org/10.1016/j.ecolmodel.2012.03.001>

979 Cristescu, M. E. (2019). Can environmental RNA revolutionize biodiversity science? *Trends in*
 980 *Ecology and Evolution*, 34(8), 694–697. doi:10.1016/j.tree.2019.05.003

981 Cristescu, M. E., & Hebert, P. D. N. N. (2018). Uses and misuses of environmental DNA in
 982 biodiversity science and conservation. *Annual Review of Ecology, Evolution, and*
 983 *Systematics*, 49(1), 209–230. doi:10.1146/annurev-ecolsys-110617-062306

984 Daly, A. J., Baetens, J. M., & De Baets, B. (2018). Ecological diversity: Measuring the
 985 unmeasurable. *Mathematics*, 6(7), 119. doi:10.3390/math6070119

986 Davies, N., Deck, J., Willis, K., Field, D., Meyer, C., Gilbert, J. A., ... Assunta-Sansone, S. (2012). A
 987 call for an international network of genomic observatories (GOs). *GigaScience*, 1(5), 1.
 988 doi:10.1186/2047-217X-1-5

989 de Menezes, A., Clipson, N., & Doyle, E. (2012). Comparative metatranscriptomics reveals
 990 widespread community responses during phenanthrene degradation in soil. *Environmental*
 991 *Microbiology*, 14(9), 2577–2588. doi:10.1111/j.1462-2920.2012.02781.x

992 Deiner, K., Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., ...
 993 Bernatchez, L. (2017). Environmental DNA metabarcoding: Transforming how we survey
 994 animal and plant communities. *Molecular Ecology*, 26(21), 5872–5895.
 995 doi:10.1111/mec.14350

996 Deiner, K., Fronhofer, E. A., Mächler, E., Walser, J. C., & Altermatt, F. (2016). Environmental
 997 DNA reveals that rivers are conveyer belts of biodiversity information. *Nature*
 998 *Communications*, 7, 12544. doi:10.1038/ncomms12544

999 Delgado-Baquerizo, M., Maestre, F. T., Reich, P. B., Jeffries, T. C., Gaitan, J. J., Encinar, D., ...
 1000 Singh, B. K. (2016). Microbial diversity drives multifunctionality in terrestrial ecosystems.
 1001 *Nature Communications*, 7, 1–8. doi:10.1038/ncomms10541

1002 Delmas, E., Besson, M., Brice, M. H., Burkle, L. A., Dalla Riva, G. V., Fortin, M. J., ... Poisot, T.
 1003 (2019). Analysing ecological networks of species interactions. *Biological Reviews*, 94(1),
 1004 16–36. doi:10.1111/brv.12433

1005 Deshpande, Reed, Sullivan, Kerkhof, Beigel, & Wade. (2019). Offline next generation

1006 metagenomics sequence analysis using MinION detection software (MINDS). *Genes*, 10(8),
1007 578. doi:10.3390/genes10080578

1008 DiBattista, J. D., Coker, D. J., Sinclair-Taylor, T. H., Stat, M., Berumen, M. L., & Bunce, M. (2017).
1009 Assessing the utility of eDNA as a tool to survey reef-fish communities in the Red Sea.
1010 *Coral Reefs*. doi:10.1007/s00338-017-1618-1

1011 Dickie, I. A., Boyer, S., Buckley, H. L., Duncan, R. P., Gardner, P. P., Hogg, I. D., ... Weaver, L.
1012 (2018). Towards robust and repeatable sampling methods in eDNA-based studies.
1013 *Molecular Ecology Resources*, 18(5), 940–952. doi:10.1111/1755-0998.12907

1014 Dowle, E., Pochon, X., Keeley, N., & Wood, S. (2015). Assessing the effects of salmon farming
1015 seabed enrichment using bacterial community diversity and high-throughput sequencing.
1016 *FEMS Microbiology Ecology*, 91(8), fiv089. doi:http://dx.doi.org/10.1093/femsec/fiv089

1017 Dufrêne, M., & Legendre, P. (1997). Species assemblages and indicator species: The need for a
1018 flexible asymmetrical approach. *Ecological Monographs*, 67(3), 345–366.
1019 doi:10.2307/2963459

1020 Dulbecco, R. (1986). A turning point in cancer research: sequencing the human genome.
1021 *Science*, 231(4742), 1055–1056. doi:10.1126/science.3945817

1022 Elbrecht, V., Braukmann, T. W. A., Ivanova, N. V., Prosser, S. W. J., Hajibabaei, M., Wright, M., ...
1023 Steinke, D. (2019). Validation of COI metabarcoding primers for terrestrial arthropods.
1024 *PeerJ*, 7, e7745. doi:10.7717/peerj.7745

1025 Elbrecht, V., & Leese, F. (2015). Can DNA-based ecosystem assessments quantify species
1026 abundance? Testing primer bias and biomass-sequence relationships with an innovative
1027 metabarcoding protocol. *PLoS ONE*, 10(7), 1–16. doi:10.1371/journal.pone.0130324

1028 Elbrecht, V., Peinert, B., & Leese, F. (2017). Sorting things out: Assessing effects of unequal
1029 specimen biomass on DNA metabarcoding. *Ecology and Evolution*, 7(17), 6918–6926.
1030 doi:10.1002/ece3.3192

1031 Elbrecht, V., Vamos, E. E., Meissner, K., Aroviita, J., & Leese, F. (2017). Assessing strengths and
1032 weaknesses of DNA metabarcoding-based macroinvertebrate identification for routine
1033 stream monitoring. *Methods in Ecology and Evolution*, 8(10), 1265–1275.
1034 doi:10.1111/2041-210X.12789

1035 Escalas, A., Hale, L., Voordeckers, J. W., Yang, Y., Firestone, M. K., Alvarez-Cohen, L., & Zhou, J.
1036 (2019). Microbial functional diversity: From concepts to applications. *Ecology and*
1037 *Evolution*, 9(20), 12000–12016. doi:10.1002/ece3.5670

1038 Fahner, N. A., Shokralla, S., Baird, D. J., & Hajibabaei, M. (2016). Large-scale monitoring of
1039 plants through environmental DNA metabarcoding of soil: Recovery, resolution, and
1040 annotation of four DNA markers. *PLoS ONE*, 11(6), e0157505.
1041 doi:10.1371/journal.pone.0157505

1042 Falk, N., Reid, T., Skoyles, A., Grgicak-Mannion, A., Drouillard, K., & Weisener, C. G. (2019).
1043 Microbial metatranscriptomic investigations across contaminant gradients of the Detroit
1044 River. *Science of The Total Environment*, 690, 121–131.
1045 doi:10.1016/j.scitotenv.2019.06.451

1046 Faust, K., Lima-Mendez, G., Lerat, J. S., Sathirapongsasuti, J. F., Knight, R., Huttenhower, C., ...
1047 Raes, J. (2015). Cross-biome comparison of microbial association networks. *Frontiers in*
1048 *Microbiology*, 6(OCT), 1–13. doi:10.3389/fmicb.2015.01200

1049 Faust, K., & Raes, J. (2012). Microbial interactions: from networks to models. *Nature Reviews*
1050 *Microbiology*, 10(8), 538–550. doi:10.1038/nrmicro2832

1051 Faust, K., Sathirapongsasuti, J. F., Izard, J., Segata, N., Gevers, D., Raes, J., & Huttenhower, C.
1052 (2012). Microbial co-occurrence relationships in the human microbiome. *PLoS*
1053 *Computational Biology*, 8(7), e1002606. doi:10.1371/journal.pcbi.1002606

1054 Ferrera, I., Giner, C. R., Reñé, A., Camp, J., Massana, R., Gasol, J. M., & Garcés, E. (2016).
1055 Evaluation of alternative high-throughput sequencing methodologies for the monitoring of
1056 marine picoplanktonic biodiversity based on rRNA gene amplicons. *Frontiers in Marine*
1057 *Science*, 3(AUG), 147. doi:10.3389/fmars.2016.00147

1058 Field, D., Garrity, G., Gray, T., Morrison, N., Selengut, J., Sterk, P., ... Wipat, A. (2008). The
1059 minimum information about a genome sequence (MIGS) specification. *Nature*
1060 *Biotechnology*, 26(5), 541–547. doi:10.1038/nbt1360

1061 Flynn, D. F. B., Mirotnick, N., Jain, M., Palmer, M. I., & Naeem, S. (2011). Functional and
1062 phylogenetic diversity as predictors of biodiversity--ecosystem-function relationships.
1063 *Ecology*, 92(8), 1573–1581. doi:10.1890/10-1245.1

1064 Fordyce, S. L., Ávila-Arcos, M. C., Rasmussen, M., Cappellini, E., Romero-Navarro, J. A., Wales,
 1065 N., ... Gilbert, M. T. P. (2013). Deep sequencing of RNA from ancient maize kernels. *PLoS*
 1066 *ONE*, 8(1), e50961. doi:10.1371/journal.pone.0050961
 1067 Friedman, J., & Alm, E. J. (2012). Inferring correlation networks from genomic survey data. *PLoS*
 1068 *Computational Biology*, 8(9), e1002687. doi:10.1371/journal.pcbi.1002687
 1069 Frontalini, F., Greco, M., Di Bella, L., Lejzerowicz, F., Reo, E., Caruso, A., ... Pawlowski, J. (2018).
 1070 Assessing the effect of mercury pollution on cultured benthic foraminifera community
 1071 using morphological and eDNA metabarcoding approaches. *Marine Pollution Bulletin*,
 1072 129(2), 512–524. doi:10.1016/j.marpolbul.2017.10.022
 1073 Frühe, L., Cordier, T., Dully, V., Breiner, H.-W., Lentendu, G., Pawlowski, J., ... Stoeck, T. (n.d.).
 1074 Supervised machine learning is superior to indicator value inference in monitoring the
 1075 environmental impacts of salmon aquaculture using eDNA metabarcodes. *Molecular*
 1076 *Ecology*, *In press*. doi:10.1111/mec.15434
 1077 Gagic, V., Bartomeus, I., Jonsson, T., Taylor, A., Winqvist, C., Fischer, C., ... Bommarco, R. (2015).
 1078 Functional identity and diversity of animals predict ecosystem functioning better than
 1079 species-based indices. *Proceedings of the Royal Society B: Biological Sciences*, 282(1801),
 1080 20142620. doi:10.1098/rspb.2014.2620
 1081 Galand, P. E., Lucas, S., Fagervold, S. K., Peru, E., Pruski, A. M., Vétion, G., ... Guizien, K. (2016).
 1082 Disturbance increases microbial community diversity and production in marine sediments.
 1083 *Frontiers in Microbiology*, 7(DEC), 1–11. doi:10.3389/fmicb.2016.01950
 1084 Galand, P. E., Pereira, O., Hochart, C., Auguet, J. C., & Debroas, D. (2018). A strong link between
 1085 marine microbial community composition and function challenges the idea of functional
 1086 redundancy. *ISME Journal*, 12(10), 2470–2478. doi:10.1038/s41396-018-0158-1
 1087 Galand, P. E., Salter, I., & Kalenitchenko, D. (2015). Ecosystem productivity is associated with
 1088 bacterial phylogenetic distance in surface marine waters. *Molecular Ecology*, 24(23), 5785–
 1089 5795. doi:10.1111/mec.13347
 1090 Gan, W., Gu, Y., Han, J., Li, C. X., Sun, J., & Liu, P. (2017). Chitosan-modified filter paper for
 1091 nucleic acid extraction and “in situ PCR” on a thermoplastic microchip. *Analytical*
 1092 *Chemistry*, 89(6), 3568–3575. doi:10.1021/acs.analchem.6b04882

1093 Gardham, S., Hose, G. C., Stephenson, S., & Chariton, A. A. (2014). DNA Metabarcoding Meets
 1094 Experimental Ecotoxicology: Advancing Knowledge on the Ecological Effects of Copper in
 1095 Freshwater Ecosystems. *Advances in Ecological Research*, 51, 79–104. doi:10.1016/B978-0-
 1096 08-099970-8.00007-5

1097 Gerhard, W. A., & Gunsch, C. K. (2019). Metabarcoding and machine learning analysis of
 1098 environmental DNA in ballast water arriving to hub ports. *Environment International*,
 1099 124(January), 312–319. doi:10.1016/j.envint.2018.12.038

1100 Gerhold, P., Cahill, J. F., Winter, M., Bartish, I. V., & Prinzing, A. (2015). Phylogenetic patterns
 1101 are not proxies of community assembly mechanisms (they are far better). *Functional
 1102 Ecology*, 29(5), 600–614. doi:10.1111/1365-2435.12425

1103 Gerlach, J., Samways, M., & Pryke, J. (2013). Terrestrial invertebrates as bioindicators: An
 1104 overview of available taxonomic groups. *Journal of Insect Conservation*, 17(4), 831–850.
 1105 doi:10.1007/s10841-013-9565-9

1106 Gibson, J. F., Shokralla, S., Curry, C., Baird, D. J., Monk, W. A., King, I., & Hajibabaei, M. (2015).
 1107 Large-scale biomonitoring of remote and threatened ecosystems via high-throughput
 1108 sequencing. *PLoS ONE*, 10(10), e0138432. doi:10.1371/journal.pone.0138432

1109 Gilbert, J. A., Meyer, F., Antonopoulos, D., Balaji, P., Brown, C. T., Brown, C. T., ... Stevens, R.
 1110 (2010). Meeting report: the terabase metagenomics workshop and the vision of an earth
 1111 microbiome project. *Standards in Genomic Sciences*, 3(3), 243–248.
 1112 doi:10.4056/sigs.1433550

1113 Goldberg, C. S., Turner, C. R., Deiner, K., Klymus, K. E., Thomsen, P. F., Murphy, M. A., ...
 1114 Taberlet, P. (2016). Critical considerations for the application of environmental DNA
 1115 methods to detect aquatic species. *Methods in Ecology and Evolution*, 7(11), 1299–1307.
 1116 doi:10.1111/2041-210X.12595

1117 Graham, E. B., Knelman, J. E., Schindlbacher, A., Siciliano, S., Breulmann, M., Yannarell, A., ...
 1118 Nemergut, D. R. (2016). Microbes as engines of ecosystem function: When does
 1119 community structure enhance predictions of ecosystem processes? *Frontiers in
 1120 Microbiology*, 7(FEB), 1–10. doi:10.3389/fmicb.2016.00214

1121 Graham, S. E., Chariton, A. A., & Landis, W. G. (2019). Using Bayesian networks to predict risk to

1122 estuary water quality and patterns of benthic environmental DNA in Queensland.
 1123 *Integrated Environmental Assessment and Management*, 15(1), 93–111.
 1124 doi:10.1002/ieam.4091

1125 Gray, C., Baird, D. J., Baumgartner, S., Jacob, U., Jenkins, G. B., O’Gorman, E. J., ... Woodward, G.
 1126 (2014). FORUM: Ecological networks: The missing links in biomonitoring science. *Journal of*
 1127 *Applied Ecology*, 51(5), 1444–1449. doi:10.1111/1365-2664.12300

1128 Grey, E. K., Bernatchez, L., Cassey, P., Deiner, K., Deveney, M., Howland, K. L., ... Lodge, D. M.
 1129 (2018). Effects of sampling effort on biodiversity patterns estimated from environmental
 1130 DNA metabarcoding surveys. *Scientific Reports*, 8(May), 8843. doi:10.1038/s41598-018-
 1131 27048-2

1132 Grizzetti, B., Lanzaova, D., Lique, C., Reynaud, A., & Cardoso, A. C. (2016). Assessing water
 1133 ecosystem services for water resource management. *Environmental Science and Policy*, 61,
 1134 194–203. doi:10.1016/j.envsci.2016.04.008

1135 Guardiola, M., Wangenstein, O. S., Taberlet, P., Coissac, E., Uriz, M. J., & Turon, X. (2016).
 1136 Spatio-temporal monitoring of deep-sea communities using metabarcoding of sediment
 1137 DNA and RNA. *PeerJ*, 4, e2807. doi:10.7717/peerj.2807

1138 Guidi, L., Chaffron, S., Bittner, L., Eveillard, D., Larhlmi, A., Roux, S., ... Gorsky, G. (2016).
 1139 Plankton networks driving carbon export in the oligotrophic ocean. *Nature*, 532(7600),
 1140 465–470. doi:10.1038/nature16942

1141 Guillou, L., Bachar, D., Audic, S., Bass, D., Berney, C., Bittner, L., ... Christen, R. (2013). The
 1142 Protist Ribosomal reference database (PR2): a catalog of unicellular eukaryote Small Sub-
 1143 Unit rRNA sequences with curated taxonomy. *Nucleic Acids Research*, 41(D1), D597–D604.
 1144 doi:10.1093/nar/gks1160

1145 Hajibabaei, M., Shokralla, S., Zhou, X., Singer, G. A. C., & Baird, D. J. (2011). Environmental
 1146 barcoding: A next-generation sequencing approach for biomonitoring applications using
 1147 river benthos. *PLoS ONE*, 6(4), e17497. doi:10.1371/journal.pone.0017497

1148 Hajibabaei, M., Spall, J. L., Shokralla, S., & van Konynenburg, S. (2012). Assessing biodiversity of
 1149 a freshwater benthic macroinvertebrate community through non-destructive
 1150 environmental barcoding of DNA from preservative ethanol. *BMC Ecology*, 12(1), 28.

doi:10.1186/1472-6785-12-28

Hauptmann, A. L., Sicheritz-Pontén, T., Cameron, K. A., Bælum, J., Plichta, D. R., Dalgaard, M., & Stibal, M. (2017). Contamination of the Arctic reflected in microbial metagenomes from the Greenland ice sheet. *Environmental Research Letters*, 12(7), 074019. doi:10.1088/1748-9326/aa7445

He, X., Sutherland, T. F., Pawlowski, J., & Abbott, C. L. (2019). Responses of foraminifera communities to aquaculture-derived organic enrichment as revealed by environmental DNA metabarcoding. *Molecular Ecology*, 28(5), 1138–1153. doi:10.1111/mec.15007

He, Z., Zhang, P., Wu, L., Rocha, A. M., Tu, Q., Shi, Z., ... Zhou, J. (2018). Microbial functional gene diversity predicts groundwater contamination and ecosystem functioning. *MBio*, 9(1), 1–15. doi:10.1128/mBio.02435-17

Hector, A., & Bagchi, R. (2007). Biodiversity and ecosystem multifunctionality. *Nature*, 448(7150), 188–190. doi:10.1038/nature05947

Hemme, C. L., Tu, Q., Shi, Z., Qin, Y., Gao, W., Deng, Y., ... Zhou, J. (2015). Comparative metagenomics reveals impact of contaminants on groundwater microbiomes. *Frontiers in Microbiology*, 6(OCT), 1–12. doi:10.3389/fmicb.2015.01205

Hering, D., Borja, A., Jones, J. I., Pont, D., Boets, P., Bouchez, A., ... Kelly, M. (2018). Implementation options for DNA-based identification into ecological status assessment under the European Water Framework Directive. *Water Research*, 138, 192–205. doi:10.1016/j.watres.2018.03.003

Hodkinson, I. D., & Jackson, J. K. (2005). Terrestrial and aquatic invertebrates as bioindicators for environmental monitoring, with particular reference to mountain ecosystems. *Environmental Management*, 35(5), 649–666. doi:10.1007/s00267-004-0211-x

Holman, L. E., de Bruyn, M., Creer, S., Carvalho, G., Robidart, J., & Rius, M. (2019). Detection of introduced and resident marine species using environmental DNA metabarcoding of sediment and water. *Scientific Reports*, 9(1), 11559. doi:10.1038/s41598-019-47899-7

Hooper Chapin III, F.S., Ewel, J.J., Hector, A., Inchausti, P., Lavorel, S., Lawton, J.H., Lodge, D.M., Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A.J., Vandermeer, J., Wardle, D.A., D. U. (2005). Effects of biodiversity on ecosystem functioning: a consensus of current

1180 knowledge. *Ecological Monographs* 2005, 75((1)), 3–35.

1181 Hughes, T. P., Kerry, J. T., Baird, A. H., Connolly, S. R., Dietzel, A., Eakin, C. M., ... Torda, G.

1182 (2018). Global warming transforms coral reef assemblages. *Nature*, 556(7702), 492–496.

1183 doi:10.1038/s41586-018-0041-2

1184 IPBES. (2019). *Summary for policymakers of the global assessment report on biodiversity and*

1185 *ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and*

1186 *Ecosystem Services. IPBES Global Assessment Summary for Policymakers* (Vol. May 2019).

1187 doi:/10.1590/1676-0611201600010001

1188 Ives, A. R., & Carpenter, S. R. (2007). Stability and diversity of ecosystems. *Science*, 317(5834),

1189 58–62. doi:10.1126/science.1133258

1190 Jarman, S. N., Berry, O., & Bunce, M. (2018). The value of environmental DNA biobanking for

1191 long-term biomonitoring. *Nature Ecology and Evolution*, 2(8), 1192–1193.

1192 doi:10.1038/s41559-018-0614-3

1193 Jeunen, G. J., Knapp, M., Spencer, H. G., Taylor, H. R., Lamare, M. D., Stat, M., ... Gemmell, N. J.

1194 (2019). Species-level biodiversity assessment using marine environmental DNA

1195 metabarcoding requires protocol optimization and standardization. *Ecology and Evolution*,

1196 9(3), 1323–1335. doi:10.1002/ece3.4843

1197 Ji, Y., Ashton, L., Pedley, S. M., Edwards, D. P., Tang, Y., Nakamura, A., ... Yu, D. W. (2013).

1198 Reliable, verifiable and efficient monitoring of biodiversity via metabarcoding. *Ecology*

1199 *Letters*, 16(10), 1245–1257. doi:10.1111/ele.12162

1200 Ji, Y., Huotari, T., Roslin, T., Schmidt, N. M., Wang, J., Yu, D. W., & Ovaskainen, O. (2019).

1201 SPIKEPIPE: A metagenomic pipeline for the accurate quantification of eukaryotic species

1202 occurrences and intraspecific abundance change using DNA barcodes or mitogenomes.

1203 *Molecular Ecology Resources*, 00, 1–12. doi:10.1111/1755-0998.13057

1204 Juul, S., Izquierdo, F., Hurst, A., Dai, X., Wright, A., Kulesha, E., ... Turner, D. J. (2015). What's in

1205 my pot? Real-time species identification on the MinION. *BioRxiv*, 030742.

1206 doi:10.1101/030742

1207 Karimi, B., Maron, P. A., Chemidlin-Prevost Boure, N., Bernard, N., Gilbert, D., & Ranjard, L.

1208 (2017). Microbial diversity and ecological networks as indicators of environmental quality.

1209 *Environmental Chemistry Letters*, 15(2), 265–281. doi:10.1007/s10311-017-0614-6
 1210 Karimi, B., Meyer, C., Gilbert, D., & Bernard, N. (2016). Air pollution below WHO levels
 1211 decreases by 40 % the links of terrestrial microbial networks. *Environmental Chemistry*
 1212 *Letters*, 14(4), 467–475. doi:10.1007/s10311-016-0589-8
 1213 Karr, J. R. (1999). Defining and measuring river health. *Freshwater Biology*, 41, 221–234.
 1214 Karsenti, E., Acinas, S. G., Bork, P., Bowler, C., De Vargas, C., Raes, J., ... Wincker, P. (2011). A
 1215 holistic approach to marine eco-systems biology. *PLoS Biology*, 9(10), e1001177.
 1216 doi:10.1371/journal.pbio.1001177
 1217 Keck, F., & Kahlert, M. (2019). Community phylogenetic structure reveals the imprint of
 1218 dispersal-related dynamics and environmental filtering by nutrient availability in
 1219 freshwater diatoms. *Scientific Reports*, 9(1), 11590. doi:10.1038/s41598-019-48125-0
 1220 Keeley, N., Wood, S. A., & Pochon, X. (2018). Development and preliminary validation of a
 1221 multi-trophic metabarcoding biotic index for monitoring benthic organic enrichment.
 1222 *Ecological Indicators*, 85, 1044–1057. doi:10.1016/j.ecolind.2017.11.014
 1223 Kelly, M., Boonham, N., Juggins, S., Kille, P., Mann, D. G., Pass, D., ... Glover, R. (2018). A DNA
 1224 based diatom metabarcoding approach for Water Framework Directive classification of
 1225 rivers. *Environment Agency, Horizon House, Deanery Road, Bristol, BS1 5AH*, (March), 157.
 1226 Kelly, M. G., Penny, C. J., & Whitton, B. A. (1995). Comparative performance of benthic diatom
 1227 indices used to assess river water quality. *Hydrobiologia*, 302(3), 179–188.
 1228 doi:10.1007/BF00032108
 1229 Kelly, R. P., Port, J. A., Yamahara, K. M., Martone, R. G., Lowell, N., Thomsen, P. F., ... Crowder, L.
 1230 B. (2014). Harnessing DNA to improve environmental management. *Science*.
 1231 doi:10.1126/science.1251156
 1232 Kermarrec, L., Franc, A., Rimet, F., Chaumeil, P., Humbert, J. F., & Bouchez, A. (2013). Next-
 1233 generation sequencing to inventory taxonomic diversity in eukaryotic communities: A test
 1234 for freshwater diatoms. *Molecular Ecology Resources*, 13(4), 607–619. doi:10.1111/1755-
 1235 0998.12105
 1236 Kermarrec, Lenaïg, Franc, A., Rimet, F., Chaumeil, P., Frigerio, J.-M., Humbert, J.-F., & Bouchez,
 1237 A. (2014). A next-generation sequencing approach to river biomonitoring using benthic

1238 diatoms. *Freshwater Science*, 33(1), 349–363. doi:10.1086/675079

1239 Kisand, V., Valente, A., Lahm, A., Tanet, G., & Lettieri, T. (2012). Phylogenetic and functional
1240 metagenomic profiling for assessing microbial biodiversity in environmental monitoring.
1241 *PLoS ONE*, 7(8), e43630. doi:10.1371/journal.pone.0043630

1242 Klunder, L., Lavaleye, M. S. S., Filippidi, A., van Bleijswijk, J. D. L., Reichart, G.-J., van der Veer, H.
1243 W., ... Mienis, F. (2018). Impact of an artificial structure on the benthic community
1244 composition in the southern North Sea: assessed by a morphological and molecular
1245 approach. *ICES Journal of Marine Science*, fsy114. doi:10.1093/icesjms/fsy114

1246 Knight, R., Vrbanac, A., Taylor, B. C., Aksenov, A., Callewaert, C., Debelius, J., ... Dorrestein, P. C.
1247 (2018). Best practices for analysing microbiomes. *Nature Reviews Microbiology*, 16(7),
1248 410–422. doi:10.1038/s41579-018-0029-9

1249 Kozlov, A. M., Zhang, J., Yilmaz, P., Glöckner, F. O., & Stamatakis, A. (2016). Phylogeny-aware
1250 identification and correction of taxonomically mislabeled sequences. *Nucleic Acids*
1251 *Research*, 44(11), 5022–5033. doi:10.1093/nar/gkw396

1252 Krehenwinkel, H., Wolf, M., Lim, J. Y., Rominger, A. J., Simison, W. B., & Gillespie, R. G. (2017).
1253 Estimating and mitigating amplification bias in qualitative and quantitative arthropod
1254 metabarcoding. *Scientific Reports*, 7(1), 1–12. doi:10.1038/s41598-017-17333-x

1255 Kurtz, Z. D., Müller, C. L., Miraldi, E. R., Littman, D. R., Blaser, M. J., & Bonneau, R. A. (2015).
1256 Sparse and Compositionally Robust Inference of Microbial Ecological Networks. *PLoS*
1257 *Computational Biology*, 11(5), 1–25. doi:10.1371/journal.pcbi.1004226

1258 Lacoursière-Roussel, A., Howland, K., Normandeau, E., Grey, E. K., Archambault, P., Deiner, K., ...
1259 Bernatchez, L. (2018). EDNA metabarcoding as a new surveillance approach for coastal
1260 Arctic biodiversity. *International Journal of Business Innovation and Research*, 17(3), 7763–
1261 7777. doi:10.1002/ece3.4213

1262 Lajoie, G., & Kembel, S. W. (2019). Making the most of trait-based approaches for microbial
1263 ecology. *Trends in Microbiology*, 27(10), 814–823. doi:10.1016/j.tim.2019.06.003

1264 Lallias, D., Hiddink, J. G., Fonseca, V. G., Gaspar, J. M., Sung, W., Neill, S. P., ... Creer, S. (2015a).
1265 Environmental metabarcoding reveals heterogeneous drivers of microbial eukaryote
1266 diversity in contrasting estuarine ecosystems. *ISME Journal*, 9(5), 1208–1221.

doi:10.1038/ismej.2014.213

Lallias, D., Hiddink, J. G., Fonseca, V. G., Gaspar, J. M., Sung, W., Neill, S. P., ... Creer, S. (2015b). Environmental metabarcoding reveals heterogeneous drivers of microbial eukaryote diversity in contrasting estuarine ecosystems. *ISME Journal*, 9(5), 1208–1221. doi:10.1038/ismej.2014.213

Langille, M. G. I., Zaneveld, J., Caporaso, J. G., McDonald, D., Knights, D., Reyes, J. A., ... Huttenhower, C. (2013). Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences. *Nature Biotechnology*, 31(9), 814–821. doi:10.1038/nbt.2676

Lanzén, A., Lekang, K., Jonassen, I., Thompson, E. M., & Troedsson, C. (2016). High-throughput metabarcoding of eukaryotic diversity for environmental monitoring of offshore oil-drilling activities. *Molecular Ecology*, 25(17), 4392–4406. doi:10.1111/mec.13761

Lanzén, A., Lekang, K., Jonassen, I., Thompson, E. M., & Troedsson, C. (2017). DNA extraction replicates improve diversity and compositional dissimilarity in metabarcoding of eukaryotes in marine sediments. *PLoS ONE*, 12(6), 1–18. doi:10.1371/journal.pone.0179443

Laroche, O., Pochon, X., Tremblay, L. A., Ellis, J. I., Lear, G., & Wood, S. A. (2018). Incorporating molecular-based functional and co-occurrence network properties into benthic marine impact assessments. *FEMS Microbiology Ecology*, 94(11), 167. doi:10.1093/femsec/fiy167

Laroche, O., Wood, S. A., Tremblay, L. A., Ellis, J. I., Lear, G., & Pochon, X. (2018). A cross-taxa study using environmental DNA/RNA metabarcoding to measure biological impacts of offshore oil and gas drilling and production operations. *Marine Pollution Bulletin*, 127, 97–107. doi:10.1016/j.marpolbul.2017.11.042

Laroche, O., Wood, S. A., Tremblay, L. A., Ellis, J. I., Lejzerowicz, F., Pawlowski, J., ... Pochon, X. (2016). First evaluation of foraminiferal metabarcoding for monitoring environmental impact from an offshore oil drilling site. *Marine Environmental Research*, 120, 225–235. doi:10.1016/j.marenvres.2016.08.009

Lau, M. K., Borrett, S. R., Baiser, B., Gotelli, N. J., & Ellison, A. M. (2017). Ecological network metrics: Opportunities for synthesis. *Ecosphere*, 8(8), :e01900. doi:10.1002/ecs2.1900

1296 Lawes, J. C., Dafforn, K. A., Clark, G. F., Brown, M. V., & Johnston, E. L. (2017). Multiple stressors
 1297 in sediments impact adjacent hard substrate habitats and across biological domains.
 1298 *Science of the Total Environment*, 592, 295–305. doi:10.1016/j.scitotenv.2017.03.083
 1299 Layeghifard, M., Hwang, D. M., & Guttman, D. S. (2017). Disentangling Interactions in the
 1300 Microbiome: A Network Perspective. *Trends in Microbiology*, 25(3), 217–228.
 1301 doi:10.1016/j.tim.2016.11.008
 1302 Lear, G., Dopheide, A., Ancion, P., & Lewis, G. D. (2011). A comparison of bacterial, ciliate and
 1303 macroinvertebrate indicators of stream ecological health. *Aquatic Ecology*, 45(4), 517–527.
 1304 doi:10.1007/s10452-011-9372-x
 1305 Leese, F., Altermatt, F., Bouchez, A., Ekrem, T., Hering, D., Meissner, K., ... Zimmermann, J.
 1306 (2016). DNAqua-Net: Developing new genetic tools for bioassessment and monitoring of
 1307 aquatic ecosystems in Europe. *Research Ideas and Outcomes*, 2, e11321.
 1308 doi:10.3897/rio.2.e11321
 1309 Leese, F., Bouchez, A., Abarenkov, K., Altermatt, F., Borja, Á., Bruce, K., ... Weigand, A. M.
 1310 (2018). Why We Need Sustainable Networks Bridging Countries, Disciplines, Cultures and
 1311 Generations for Aquatic Biomonitoring 2.0: A Perspective Derived From the DNAqua-Net
 1312 COST Action. *Advances in Ecological Research*, in press, 63–99.
 1313 doi:10.1016/bs.aecr.2018.01.001
 1314 Lejzerowicz, F., Esling, P., Pillet, L., Wilding, T. A., Black, K. D., & Pawlowski, J. (2015). High-
 1315 throughput sequencing and morphology perform equally well for benthic monitoring of
 1316 marine ecosystems. *Scientific Reports*, 5(1), 13932. doi:10.1038/srep13932
 1317 Leray, M., Knowlton, N., Ho, S.-L., Nguyen, B. N., & Machida, R. J. (2019). GenBank is a reliable
 1318 resource for 21st century biodiversity research. *Proceedings of the National Academy of*
 1319 *Sciences*, 201911714. doi:10.1073/pnas.1911714116
 1320 Levin, L. A., Bett, B. J., Gates, A. R., Heimbach, P., Howe, B. M., Janssen, F., ... Weller, R. A.
 1321 (2019). Global observing needs in the deep ocean. *Frontiers in Marine Science*, 6(May),
 1322 241. doi:10.3389/fmars.2019.241
 1323 Li, F., Peng, Y., Fang, W., Altermatt, F., Xie, Y., Yang, J., & Zhang, X. (2018). Application of
 1324 environmental DNA metabarcoding for predicting anthropogenic pollution in rivers.

1325 *Environmental Science and Technology*, 52(20), 11708–11719.
 1326 doi:10.1021/acs.est.8b03869

1327 Libbrecht, M. W., & Noble, W. S. (2015). Machine learning applications in genetics and
 1328 genomics. *Nat Rev Genet*, 16(6), 321–332. doi:10.1038/nrg3920

1329 Lima-Mendez, G., Faust, K., Henry, N., Decelle, J., Colin, S., Carcillo, F., ... Velayoudon, D. (2015).
 1330 Determinants of community structure in the global plankton interactome. *Science*,
 1331 348(6237). doi:10.1126/science.1262073

1332 Liu, C., Yao, M., Stegen, J. C., Rui, J., Li, J., & Li, X. (2017). Long-term nitrogen addition affects the
 1333 phylogenetic turnover of soil microbial community responding to moisture pulse. *Scientific*
 1334 *Reports*, 7(1), 17492. doi:10.1038/s41598-017-17736-w

1335 Lobo, J., Shokralla, S., Costa, M. H., Hajibabaei, M., & Costa, F. O. (2017). DNA metabarcoding
 1336 for high-throughput monitoring of estuarine macrobenthic communities. *Scientific*
 1337 *Reports*, 7(1), 15618. doi:10.1038/s41598-017-15823-6

1338 Loreau, M., & de Mazancourt, C. (2013). Biodiversity and ecosystem stability: A synthesis of
 1339 underlying mechanisms. *Ecology Letters*, 16, 106–115. doi:10.1111/ele.12073

1340 Louca, S., Jacques, S. M. S., Pires, A. P. F., Leal, J. S., Srivastava, D. S., Parfrey, L. W., ... Doebeli,
 1341 M. (2016). High taxonomic variability despite stable functional structure across microbial
 1342 communities. *Nature Ecology & Evolution*, 1, 0015. doi:10.1038/s41559-016-0015

1343 Louca, S., Parfrey, L. W., & Doebeli, M. (2016). Decoupling function and taxonomy in the global
 1344 ocean microbiome. *Science*, 353(6305), 1272–1277. doi:10.1126/science.aaf4507

1345 Louca, S., Polz, M. F., Mazel, F., Albright, M. B. N., Huber, J. A., O'Connor, M. I., ... Parfrey, L. W.
 1346 (2018). Function and functional redundancy in microbial systems. *Nature Ecology and*
 1347 *Evolution*, 2(6), 936–943. doi:10.1038/s41559-018-0519-1

1348 Lupatini, M., Suleiman, A. K. A., Jacques, R. J. S., Antonioli, Z. I., de Siqueira Ferreira, A.,
 1349 Kuramae, E. E., & Roesch, L. F. W. (2014). Network topology reveals high connectance
 1350 levels and few key microbial genera within soils. *Frontiers in Environmental Science*,
 1351 2(May), 1–11. doi:10.3389/fenvs.2014.00010

1352 Ma, A., Lu, X., Gray, C., Raybould, A., Tamaddon-Nezhad, A., Woodward, G., & Bohan, D. A.
 1353 (2019). Ecological networks reveal resilience of agro-ecosystems to changes in farming

1354 management. *Nature Ecology and Evolution*, 3(2), 260–264. doi:10.1038/s41559-018-
 1355 0757-2

1356 Manning, P., Van Der Plas, F., Soliveres, S., Allan, E., Maestre, F. T., Mace, G., ... Fischer, M.
 1357 (2018). Redefining ecosystem multifunctionality. *Nature Ecology and Evolution*, 2(3), 427–
 1358 436. doi:10.1038/s41559-017-0461-7

1359 Manolio, T. A., Chisholm, R. L., Ozenberger, B., Roden, D. M., Williams, M. S., Wilson, R., ...
 1360 Ginsburg, G. S. (2013). Implementing genomic medicine in the clinic: The future is here.
 1361 *Genetics in Medicine*, 15(4), 258–267. doi:10.1038/gim.2012.157

1362 Martínez-Santos, M., Lanzén, A., Unda-Calvo, J., Martín, I., Garbisu, C., & Ruiz-Romera, E.
 1363 (2018). Treated and untreated wastewater effluents alter river sediment bacterial
 1364 communities involved in nitrogen and sulphur cycling. *Science of the Total Environment*,
 1365 633, 1051–1061. doi:10.1016/j.scitotenv.2018.03.229

1366 Martiny, A. C., Treseder, K., & Pusch, G. (2013). Phylogenetic conservatism of functional traits in
 1367 microorganisms. *The ISME Journal*, 7(4), 830–8. doi:10.1038/ismej.2012.160

1368 Marull, J., Pino, J., Mallarach, J. M., & Cordobilla, M. J. (2007). A land suitability index for
 1369 strategic environmental assessment in metropolitan areas. *Landscape and Urban Planning*,
 1370 81(3), 200–212. doi:10.1016/j.landurbplan.2006.11.005

1371 Maurer, D. O. N., Nguyen, H. A. I., Robertson, G., & Gerlinger, T. O. M. (1999). The Infaunal
 1372 Trophic Index (ITI): Its suitability for marine environmental monitoring. *Ecological*
 1373 *Applications*, 9(June 1997), 699–713. doi:10.1890/1051-
 1374 0761(1999)009[0699:TITIII]2.0.CO;2

1375 Mayfield, M. M., & Levine, J. M. (2010). Opposing effects of competitive exclusion on the
 1376 phylogenetic structure of communities. *Ecology Letters*, 13(9), 1085–1093.
 1377 doi:10.1111/j.1461-0248.2010.01509.x

1378 Mazel, F., Pennell, M. W., Cadotte, M. W., Diaz, S., Dalla Riva, G. V., Grenyer, R., ... Pearse, W. D.
 1379 (2018). Prioritizing phylogenetic diversity captures functional diversity unreliably. *Nature*
 1380 *Communications*, 9(1), 2888. doi:10.1038/s41467-018-05126-3

1381 McCann, K. S. (2000). The diversity–stability debate. *Nature*, 405(May), 228–233.

1382 McDonald, D., Clemente, J. C., Kuczynski, J., Rideout, J. R., Stombaugh, J., Wendel, D., ...

1383 Caporaso, J. G. (2012). The Biological Observation Matrix (BIOM) format or: How I learned
 1384 to stop worrying and love the ome-ome. *GigaScience*, 1(1), 464. doi:10.1186/2047-217X-1-
 1385 7

1386 McFarlin, K. M., Questel, J. M., Hopcroft, R. R., & Leigh, M. B. (2017). Bacterial community
 1387 structure and functional potential in the northeastern Chukchi Sea. *Continental Shelf*
 1388 *Research*, 136, 20–28. doi:10.1016/j.csr.2017.01.018

1389 McGee, K. M., Robinson, C. V., & Hajibabaei, M. (2019). Gaps in DNA-Based Biomonitoring
 1390 Across the Globe. *Frontiers in Ecology and Evolution*, 7, 337. doi:10.3389/fevo.2019.00337

1391 Momal, R., Robin, S., & Ambroise, C. (2019). Tree-based reconstruction of ecological network
 1392 from abundance data. *ArXiv*, 1–23. Retrieved from <http://arxiv.org/abs/1905.02452>

1393 Morris, Z. S., wooding, S., & Grant, J. (2011). The answer is 17 years, what is the question:
 1394 Understanding time lags in translational research. *Journal of the Royal Society of Medicine*,
 1395 104(12), 510–520. doi:10.1258/jrsm.2011.110180

1396 Mortágua, A., Vasselon, V., Oliveira, R., Elias, C., Chardon, C., Bouchez, A., ... F.P. Almeida, S.
 1397 (2019). Applicability of DNA metabarcoding approach in the bioassessment of Portuguese
 1398 rivers using diatoms. *Ecological Indicators*, 106, 105470.
 1399 doi:10.1016/j.ecolind.2019.105470

1400 Mouillot, D., Graham, N. A. J., Villéger, S., Mason, N. W. H., & Bellwood, D. R. (2013). A
 1401 functional approach reveals community responses to disturbances. *Trends in Ecology and*
 1402 *Evolution*, 28(3), 167–177. doi:10.1016/j.tree.2012.10.004

1403 Mukherjee, A., Chettri, B., Langpoklakpam, J. S., Basak, P., Prasad, A., Mukherjee, A. K., ...
 1404 Chattopadhyay, D. (2017). Bioinformatic approaches including predictive metagenomic
 1405 profiling reveal characteristics of bacterial response to petroleum hydrocarbon
 1406 contamination in diverse environments. *Scientific Reports*, 7(1), 1108. doi:10.1038/s41598-
 1407 017-01126-3

1408 Nascimento, F. J. A., Lallias, D., Bik, H. M., & Creer, S. (2018). Sample size effects on the
 1409 assessment of eukaryotic diversity and community structure in aquatic sediments using
 1410 high-throughput sequencing. *Scientific Reports*, 8(1), 11737. doi:10.1038/s41598-018-
 1411 30179-1

1412 Nguyen, T. T., Cochrane, S. K. J., & Landfald, B. (2018). Perturbation of seafloor bacterial
 1413 community structure by drilling waste discharge. *Marine Pollution Bulletin*, 129(2), 615–
 1414 622. doi:10.1016/j.marpolbul.2017.10.039
 1415 Niemeijer, D. (2002). Developing indicators for environmental policy: Data-driven and theory-
 1416 driven approaches examined by example. *Environmental Science and Policy*, 5(2), 91–103.
 1417 doi:10.1016/S1462-9011(02)00026-6
 1418 Nilsson, L. K. J., Sharma, A., Bhatnagar, R. K., Bertilsson, S., & Terenius, O. (2018). Presence of
 1419 Aedes and Anopheles mosquito larvae is correlated to bacteria found in domestic water-
 1420 storage containers. *FEMS Microbiology Ecology*, 94(6), fiy058. doi:10.1093/femsec/fiy058
 1421 Nilsson, R. H., Larsson, K. H., Taylor, A. F. S., Bengtsson-Palme, J., Jeppesen, T. S., Schigel, D., ...
 1422 Abarenkov, K. (2019). The UNITE database for molecular identification of fungi: Handling
 1423 dark taxa and parallel taxonomic classifications. *Nucleic Acids Research*, 47(D1), D259–
 1424 D264. doi:10.1093/nar/gky1022
 1425 Obi, C. C., Adebuseye, S. A., Ugoji, E. O., Ilori, M. O., Amund, O. O., & Hickey, W. J. (2016).
 1426 Microbial communities in sediments of Lagos Lagoon, Nigeria: Elucidation of community
 1427 structure and potential impacts of contamination by municipal and industrial wastes.
 1428 *Frontiers in Microbiology*, 7(AUG), 1213. doi:10.3389/fmicb.2016.01213
 1429 Oliver, T. H., Heard, M. S., Isaac, N. J. B., Roy, D. B., Procter, D., Eigenbrod, F., ... Bullock, J. M.
 1430 (2015). Biodiversity and resilience of ecosystem functions. *Trends in Ecology and Evolution*,
 1431 30(11), 673–684. doi:10.1016/j.tree.2015.08.009
 1432 Pagenkopp Lohan, K. M., Fleischer, R. C., Carney, K. J., Holzer, K. K., & Ruiz, G. M. (2016).
 1433 Amplicon-based pyrosequencing reveals high diversity of protistan parasites in ships'
 1434 ballast water: implications for biogeography and infectious diseases. *Microbial Ecology*,
 1435 71(3), 530–542. doi:10.1007/s00248-015-0684-6
 1436 Pauvert, C., Vallance, J., Delière, L., Buée, M., & Vacher, C. (2019). Microbial networks inferred
 1437 from metabarcoding data lack replicability: consequences for next-generation
 1438 biomonitoring. *BioRxiv*, 642199. doi:https://doi.org/10.1101/642199
 1439 Pawlowski, J., Lejzerowicz, F., Apotheloz-Perret-Gentil, L., Visco, J., & Esling, P. (2016). Protist
 1440 metabarcoding and environmental biomonitoring: Time for change. *European Journal of*

1441 *Protistology*, 55, Part A, 12–25. doi:http://dx.doi.org/10.1016/j.ejop.2016.02.003

1442 Pawlowski, Jan, Esling, P., Lejzerowicz, F., Cedhagen, T., & Wilding, T. A. (2014). Environmental
 1443 monitoring through protist next-generation sequencing metabarcoding: Assessing the
 1444 impact of fish farming on benthic foraminifera communities. *Molecular Ecology Resources*,
 1445 14(6), 1129–1140. doi:10.1111/1755-0998.12261

1446 Pawlowski, Jan, Esling, P., Lejzerowicz, F., Cordier, T., Visco, J. A., Martins, C. I. M., ... Cedhagen,
 1447 T. (2016). Benthic monitoring of salmon farms in Norway using foraminiferal
 1448 metabarcoding. *Aquaculture Environment Interactions*, 8, 371–386.

1449 Pawlowski, Jan, Kelly-Quinn, M., Altermatt, F., Apothéloz-Perret-Gentil, L., Beja, P., Boggero, A.,
 1450 ... Kahlert, M. (2018). The future of biotic indices in the ecogenomic era: Integrating
 1451 (e)DNA metabarcoding in biological assessment of aquatic ecosystems. *Science of the Total*
 1452 *Environment*, 637–638, 1295–1310. doi:10.1016/j.scitotenv.2018.05.002

1453 Pawlowski, Jan, Lejzerowicz, F., & Esling, P. (2014). Next-generation environmental diversity
 1454 surveys of foraminifera: Preparing the future. *Biological Bulletin*, 227(2), 93–106.

1455 Payne, R. J. (2013). Seven reasons why protists make useful bioindicators. *Acta Protozoologica*,
 1456 52(3), 105–113. doi:10.4467/16890027AP.13.0011.1108

1457 Pearman, J. K., Aylagas, E., Voolstra, C. R., Anlauf, H., Villalobos, R., & Carvalho, S. (2019).
 1458 Disentangling the complex microbial community of coral reefs using standardized
 1459 Autonomous Reef Monitoring Structures (ARMS). *Molecular Ecology*, 28(15), 3496–3507.
 1460 doi:10.1111/mec.15167

1461 Pellissier, L., Albouy, C., Bascompte, J., Farwig, N., Graham, C., Loreau, M., ... Gravel, D. (2018).
 1462 Comparing species interaction networks along environmental gradients. *Biological*
 1463 *Reviews*, 93(2), 785–800. doi:10.1111/brv.12366

1464 Pérez-Valera, E., Goberna, M., Faust, K., Raes, J., García, C., & Verdú, M. (2017). Fire modifies
 1465 the phylogenetic structure of soil bacterial co-occurrence networks. *Environmental*
 1466 *Microbiology*, 19(1), 317–327. doi:10.1111/1462-2920.13609

1467 Piñol, J., Mir, G., Gomez-Polo, P., & Agustí, N. (2015). Universal and blocking primer mismatches
 1468 limit the use of high-throughput DNA sequencing for the quantitative metabarcoding of
 1469 arthropods. *Molecular Ecology Resources*, 15(4), 819–830. doi:10.1111/1755-0998.12355

1470 Pitsch, G., Bruni, E. P., Forster, D., Qu, Z., Sonntag, B., Stoeck, T., & Posch, T. (2019). Seasonality
 1471 of planktonic freshwater ciliates: Are analyses based on V9 regions of the 18S rRNA gene
 1472 correlated with morphospecies counts? *Frontiers in Microbiology*, *10*(FEB), 1–15.
 1473 doi:10.3389/fmicb.2019.00248

1474 Pochon, X., Wood, S. A., Keeley, N. B., Lejzerowicz, F., Esling, P., Drew, J., & Pawlowski, J. (2015).
 1475 Accurate assessment of the impact of salmon farming on benthic sediment enrichment
 1476 using foraminiferal metabarcoding. *Marine Pollution Bulletin*, *100*(1), 370–382.
 1477 doi:10.1016/j.marpolbul.2015.08.022

1478 Pochon, Xavier, Zaiko, A., Fletcher, L. M., Laroche, O., & Wood, S. A. (2017). Wanted dead or
 1479 alive? Using metabarcoding of environmental DNA and RNA to distinguish living
 1480 assemblages for biosecurity applications. *PLoS ONE*, *12*(11), 1–19.
 1481 doi:10.1371/journal.pone.0187636

1482 Poikane, S., Zampoukas, N., Borja, A., Davies, S. P., van de Bund, W., & Birk, S. (2014).
 1483 Intercalibration of aquatic ecological assessment methods in the European Union: Lessons
 1484 learned and way forward. *Environmental Science and Policy*, *44*, 237–246.
 1485 doi:10.1016/j.envsci.2014.08.006

1486 Pollock, J., Glendinning, L., Wisedchanwet, T., & Watson, M. (2018). The madness of
 1487 microbiome : attempting to find consensus. *Applied and Environmental Microbiology*,
 1488 *84*(7), 1–12.

1489 Pont, D., Hugueny, B., Beier, U., Goffaux, D., Melcher, A., Noble, R., ... Schmutz, S. (2006).
 1490 Assessing river biotic condition at a continental scale: A European approach using
 1491 functional metrics and fish assemblages. *Journal of Applied Ecology*, *43*(1), 70–80.
 1492 doi:10.1111/j.1365-2664.2005.01126.x

1493 Prygiel, J., & Coste, M. (2000). Guide méthodologique pour la mise en oeuvre de l'Indice
 1494 Biologique Diatomées NF T 90-354. *Agences de l'Eau, Ministère de l'Aménagement Du*
 1495 *Territoire et de l'Environnement, Direction de l'Eau & CEMAGREF, Paris, France.*

1496 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., ... Glöckner, F. O. (2013). The
 1497 SILVA ribosomal RNA gene database project: Improved data processing and web-based
 1498 tools. *Nucleic Acids Research*, *41*, D590–D596. doi:10.1093/nar/gks1219

1499 Quince, C., Walker, A. W., Simpson, J. T., Loman, N. J., & Segata, N. (2017). Shotgun
 1500 metagenomics, from sampling to analysis. *Nature Biotechnology*, 35(12), 1211.
 1501 doi:10.1038/nbt1217-1211b

1502 Quinn, R. A., Navas-Molina, J. A., Hyde, E. R., Song, S. J., Vázquez-Baeza, Y., Humphrey, G., ...
 1503 Knight, R. (2016). From sample to multi-omics conclusions in under 48 hours. *MSystems*,
 1504 1(2), e00038-16. doi:10.1128/msystems.00038-16

1505 Raes, J., Letunic, I., Yamada, T., Jensen, L. J., & Bork, P. (2011). Toward molecular trait-based
 1506 ecology through integration of biogeochemical, geographical and metagenomic data.
 1507 *Molecular Systems Biology*, 7(473), 1–9. doi:10.1038/msb.2011.6

1508 Ramond, P., Sourisseau, M., Simon, N., Romac, S., Schmitt, S., Rigaut-Jalabert, F., ... Siano, R.
 1509 (2019). Coupling between taxonomic and functional diversity in protistan coastal
 1510 communities. *Environmental Microbiology*, 21(2), 730–749. doi:10.1111/1462-2920.14537

1511 Ratajczak, Z., Carpenter, S. R., Ives, A. R., Kucharik, C. J., Ramiadantsoa, T., Stegner, M. A., ...
 1512 Turner, M. G. (2018). Abrupt change in ecological systems: inference and diagnosis. *Trends*
 1513 *in Ecology and Evolution*, 33(7), 513–526. doi:10.1016/j.tree.2018.04.013

1514 Ratnasingham, S., & Hebert, P. D. N. (2007). BOLD: The barcode of life data system: barcoding.
 1515 *Molecular Ecology Notes*, 7(3), 355–364. doi:10.1111/j.1471-8286.2007.01678.x

1516 Reintjes, G., Tegetmeyer, H. E., Bürgisser, M., Orlić, S., Tews, I., Zubkov, M., ... Fuchs, B. M.
 1517 (2019). On-Site analysis of bacterial communities of the ultraoligotrophic South Pacific
 1518 Gyre. *Applied and Environmental Microbiology*, 85(14). doi:10.1128/AEM.00184-19

1519 Reynoldson, T. B., & Metcalfe-Smith, J. L. (1992). An overview of the assessment of aquatic
 1520 ecosystem health using benthic invertebrates. *Journal of Aquatic Ecosystem Health*, 1(4),
 1521 295–308. doi:10.1007/BF00044171

1522 Rimet, F., Chaumeil, P., Keck, F., Kermarrec, L., Vasselon, V., Kahlert, M., ... Bouchez, A. (2016).
 1523 R-Syst::diatom: An open-access and curated barcode database for diatoms and freshwater
 1524 monitoring. *Database*, 2016:, baw016. doi:10.1093/database/baw016

1525 Rivera, S. F., Vasselon, V., Jacquet, S., Bouchez, A., Ariztegui, D., & Rimet, F. (2018).
 1526 Metabarcoding of lake benthic diatoms: from structure assemblages to ecological
 1527 assessment. *Hydrobiologia*, 807(1), 37–51. doi:10.1007/s10750-017-3381-2

1528 Rivera, Sinziana F., Vasselon, V., Bouchez, A., & Rimet, F. (2020). Diatom metabarcoding applied
 1529 to large scale monitoring networks: Optimization of bioinformatics strategies using Mothur
 1530 software. *Ecological Indicators*, 109, 105775. doi:10.1016/j.ecolind.2019.105775
 1531 Roy, S., Coldren, C., Karunamurthy, A., Kip, N. S., Klee, E. W., Lincoln, S. E., ... Carter, A. B.
 1532 (2018). Standards and guidelines for validating next-generation sequencing bioinformatics
 1533 pipelines: a joint recommendation of the association for molecular pathology and the
 1534 college of american pathologists. *Journal of Molecular Diagnostics*, 20(1), 4–27.
 1535 doi:10.1016/j.jmoldx.2017.11.003
 1536 Ruppert, K. M., Kline, R. J., & Rahman, M. S. (2019). Past, present, and future perspectives of
 1537 environmental DNA (eDNA) metabarcoding: A systematic review in methods, monitoring,
 1538 and applications of global eDNA. *Global Ecology and Conservation*, 17, e00547.
 1539 doi:10.1016/j.gecco.2019.e00547
 1540 Rygg, B., & Norling, K. (2013). *Norwegian Sensitivity Index (NSI) for marine macroinvertebrates,*
 1541 *and an update of Indicator Species Index (ISI)*. ISBN:978-82-577-6210-0.
 1542 Salis, R. K., Bruder, A., Piggott, J. J., Summerfield, T. C., & Matthaei, C. D. (2017). High-
 1543 throughput amplicon sequencing and stream benthic bacteria: Identifying the best
 1544 taxonomic level for multiple-stressor research. *Scientific Reports*, 7, 44657.
 1545 doi:10.1038/srep44657
 1546 Santini, L., Belmaker, J., Costello, M. J., Pereira, H. M., Rossberg, A. G., Schipper, A. M., ...
 1547 Rondinini, C. (2017). Assessing the suitability of diversity metrics to detect biodiversity
 1548 change. *Biological Conservation*, 213, 341–350. doi:10.1016/j.biocon.2016.08.024
 1549 Shade, A. (2017). Diversity is the question, not the answer. *ISME Journal*, 11(1), 1–6.
 1550 doi:10.1038/ismej.2016.118
 1551 Sigsgaard, E. E., Nielsen, I. B., Bach, S. S., Lorenzen, E. D., Robinson, D. P., Knudsen, S. W., ...
 1552 Thomsen, P. F. (2016). Population characteristics of a large whale shark aggregation
 1553 inferred from seawater environmental DNA. *Nature Ecology & Evolution*, 1(1), 0004.
 1554 doi:10.1038/s41559-016-0004
 1555 Simonin, M., Voss, K. A., Hassett, B. A., Rocca, J. D., Wang, S. Y., Bier, R. L., ... Bernhardt, E. S.
 1556 (2019). In search of microbial indicator taxa: shifts in stream bacterial communities along

1557 an urbanization gradient. *Environmental Microbiology*, 21(10), 3653–3668.
 1558 doi:10.1111/1462-2920.14694
 1559 Singer, E., Wagner, M., & Woyke, T. (2017). Capturing the genetic makeup of the active
 1560 microbiome in situ. *ISME Journal*, 11(9), 1949–1963. doi:10.1038/ismej.2017.59
 1561 Smith, M. B., Rocha, A. M., Smillie, C. S., Olesen, S. W., Paradis, C., Wu, L., ... Hazen, T. C. (2015).
 1562 Natural bacterial communities serve as quantitative geochemical biosensors. *MBio*, 6(3),
 1563 1–13. doi:10.1128/mBio.00326-15
 1564 Srivastava, D. S., Cadotte, M. W., Macdonald, A. A. M., Marushia, R. G., & Mirotnick, N.
 1565 (2012). Phylogenetic diversity and the functioning of ecosystems. *Ecology Letters*, 15(7),
 1566 637–648. doi:10.1111/j.1461-0248.2012.01795.x
 1567 Stark, J. D. (1998). SQMCI: A biotic index for freshwater macroinvertebrate coded-abundance
 1568 data. *New Zealand Journal of Marine and Freshwater Research*, 32(1), 55–66.
 1569 doi:10.1080/00288330.1998.9516805
 1570 Stark, Z., Dolman, L., Manolio, T. A., Ozenberger, B., Hill, S. L., Caulfield, M. J., ... North, K. N.
 1571 (2019). Integrating genomics into healthcare: a global responsibility. *American Journal of*
 1572 *Human Genetics*, 104(1), 13–20. doi:10.1016/j.ajhg.2018.11.014
 1573 Stat, M., Huggett, M. J., Bernasconi, R., Dibattista, J. D., Berry, T. E., Newman, S. J., ... Bunce, M.
 1574 (2017). Ecosystem biomonitoring with eDNA: Metabarcoding across the tree of life in a
 1575 tropical marine environment. *Scientific Reports*, 7, 12240. doi:10.1038/s41598-017-12501-
 1576 5
 1577 Stefanni, S., Stanković, D., Borme, D., de Olazabal, A., Juretić, T., Pallavicini, A., & Tirelli, V.
 1578 (2018). Multi-marker metabarcoding approach to study mesozooplankton at basin scale.
 1579 *Scientific Reports*, 8(1), 12085. doi:10.1038/s41598-018-30157-7
 1580 Stoeck, T., Frühe, L., Forster, D., Cordier, T., Martins, C. I. M., & Pawlowski, J. (2018).
 1581 Environmental DNA metabarcoding of benthic bacterial communities indicates the benthic
 1582 footprint of salmon aquaculture. *Marine Pollution Bulletin*, 127(November 2017), 139–149.
 1583 doi:10.1016/j.marpolbul.2017.11.065
 1584 Stoeck, T., Kochems, R., Forster, D., Lejzerowicz, F., & Pawlowski, J. (2018). Metabarcoding of
 1585 benthic ciliate communities shows high potential for environmental monitoring in salmon

1586 aquaculture. *Ecological Indicators*, 85, 153–164. doi:10.1016/j.ecolind.2017.10.041

1587 Stork, N. E. (2018). How many species of insects and other terrestrial arthropods are there on

1588 Earth? *Annual Review of Entomology*, 63(1), 31–45. doi:10.1146/annurev-ento-020117-

1589 043348

1590 Sun, D. L., Jiang, X., Wu, Q. L., & Zhou, N. Y. (2013). Intragenomic heterogeneity of 16S rRNA

1591 genes causes overestimation of prokaryotic diversity. *Applied and Environmental*

1592 *Microbiology*, 79(19), 5962–5969. doi:10.1128/AEM.01282-13

1593 Taberlet, P., Bonin, A., Zinger, L., & Coissac, E. (2018). *Environmental DNA: For Biodiversity*

1594 *Research and Monitoring*. Oxford University Press.

1595 doi:10.1093/oso/9780198767220.001.0001

1596 Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C., & Willerslev, E. (2012). Towards next-

1597 generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology*, 21(8),

1598 2045–2050. doi:10.1111/j.1365-294X.2012.05470.x

1599 Tackmann, J., Matias Rodrigues, J. F., & von Mering, C. (2019). Rapid inference of direct

1600 interactions in large-scale ecological networks from heterogeneous microbial sequencing

1601 data. *Cell Systems*, 9, 286–296. doi:10.1016/j.cels.2019.08.002

1602 Tamaddoni-Nezhad, A., Milani, G. A., Raybould, A., Muggleton, S., & Bohan, D. A. (2013).

1603 Construction and validation of food webs using logic-based machine learning and text

1604 mining. *Advances in Ecological Research*, 49, 225–289. doi:10.1016/B978-0-12-420002-

1605 9.00004-4

1606 Tapolczai, K., Keck, F., Bouchez, A., Rimet, F., & Vasselon, V. (2019). Diatom DNA metabarcoding

1607 for biomonitoring : strategies to avoid major taxonomical and bioinformatical biases

1608 limiting molecular indices capacities. *Frontiers in Ecology and Evolution*, 7(409). doi:doi:

1609 10.3389/fevo.2019.00409

1610 Tapolczai, K., Vasselon, V., Bouchez, A., Stenger-Kovács, C., Padisák, J., & Rimet, F. (2019). The

1611 impact of OTU sequence similarity threshold on diatom-based bioassessment: A case study

1612 of the rivers of Mayotte (France, Indian Ocean). *Ecology and Evolution*, 9(1), 166–179.

1613 doi:10.1002/ece3.4701

1614 Thompson, M. S. A., Bankier, C., Bell, T., Dumbrell, A. J., Gray, C., Ledger, M. E., ... Woodward, G.

1615 (2016). Gene-to-ecosystem impacts of a catastrophic pesticide spill: testing a multilevel
 1616 bioassessment approach in a river ecosystem. *Freshwater Biology*, 61(12), 2037–2050.
 1617 doi:10.1111/fwb.12676

1618 Thomsen, P. F., Kielgast, J., Iversen, L. L., Møller, P. R., Rasmussen, M., & Willerslev, E. (2012).
 1619 Detection of a diverse marine fish fauna using environmental DNA from seawater samples.
 1620 *PLoS ONE*, 7(8), e41732. doi:10.1371/journal.pone.0041732

1621 Tilman, D., Reich, P. B., & Knops, J. M. H. (2006). Biodiversity and ecosystem stability in a
 1622 decade-long grassland experiment. *Nature*, 441(7093), 629–632. doi:10.1038/nature04742

1623 Tkacz, A., Hortala, M., & Poole, P. S. (2018). Absolute quantitation of microbiota abundance in
 1624 environmental samples. *Microbiome*, 6(1), 1–13. doi:10.1186/s40168-018-0491-7

1625 Tucker, C. M., Cadotte, M. W., Carvalho, S. B., Jonathan Davies, T., Ferrier, S., Fritz, S. A., ...
 1626 Mazel, F. (2017). A guide to phylogenetic metrics for conservation, community ecology and
 1627 macroecology. *Biological Reviews*, 92(2), 698–715. doi:10.1111/brv.12252

1628 Tylianakis, J. M., & Morris, R. J. (2017). Ecological networks across environmental gradients.
 1629 *Annual Review of Ecology, Evolution, and Systematics*, 48(1), annurev-ecolsys-110316-
 1630 022821. doi:10.1146/annurev-ecolsys-110316-022821

1631 Tylianakis, J. M., Tscharnkte, T., & Lewis, O. T. (2007). Habitat modification alters the structure
 1632 of tropical host-parasitoid food webs. *Nature*, 445(7124), 202–205.
 1633 doi:10.1038/nature05429

1634 United Nations. (n.d.). A/RES/70/1: Sustainable Development Knowledge Platform.

1635 Urzelai, A., Hernández, A. J., & Pastor, J. (2000). Biotic indices based on soil nematode
 1636 communities for assessing soil quality in terrestrial ecosystems. *Science of the Total
 1637 Environment*, 247(2–3), 253–261. doi:10.1016/S0048-9697(99)00494-5

1638 Vacher, C., Tamaddoni-Nezhad, A., Kamenova, S., Peyrard, N., Moalic, Y., Sabbadin, R., ... Bohan,
 1639 D. A. (2016). Learning ecological networks from next-generation sequencing data. In
 1640 *Advances in Ecological Research* (Vol. 54, pp. 1–39). doi:10.1016/bs.aecr.2015.10.004

1641 Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J., Thomsen, P. F., ... Dejean, T. (2016).
 1642 Next-generation monitoring of aquatic biodiversity using environmental DNA
 1643 metabarcoding. *Molecular Ecology*, 25(4), 929–942. doi:10.1111/mec.13428

1644 Vandenkoornhuyse, P., Dufresne, A., Quaiser, A., Gouesbet, G., Binet, F., Francez, A. J., ...
 1645 Couée, I. (2010). Integration of molecular functions at the ecosystemic level:
 1646 Breakthroughs and future goals of environmental genomics and post-genomics. *Ecology*
 1647 *Letters*, 13(6), 776–791. doi:10.1111/j.1461-0248.2010.01464.x
 1648 Vasselon, V., Bouchez, A., Rimet, F., Jacquet, S., Trobajo, R., Corniquel, M., ... Domaizon, I.
 1649 (2018). Avoiding quantification bias in metabarcoding: Application of a cell biovolume
 1650 correction factor in diatom molecular biomonitoring. *Methods in Ecology and Evolution*,
 1651 9(4), 1060–1069. doi:10.1111/2041-210X.12960
 1652 Vasselon, V., Domaizon, I., Rimet, F., Kahlert, M., & Bouchez, A. (2017). Application of high-
 1653 throughput sequencing (HTS) metabarcoding to diatom biomonitoring: Do DNA extraction
 1654 methods matter? *Freshwater Science*, 36(1), 162–177. doi:10.1086/690649
 1655 Vasselon, V., Rimet, F., Tapolczai, K., & Bouchez, A. (2017). Assessing ecological status with
 1656 diatoms DNA metabarcoding: Scaling-up on a WFD monitoring network (Mayotte island,
 1657 France). *Ecological Indicators*, 82, 1–12. doi:10.1016/j.ecolind.2017.06.024
 1658 Venail, P. A., & Vives, M. J. (2013). Phylogenetic distance and species richness interactively
 1659 affect the productivity of bacterial communities. *Ecology*, 94(11), 2529–2536.
 1660 doi:10.1890/12-2002.1
 1661 Větrovský, T., & Baldrian, P. (2013). The variability of the 16S rRNA gene in bacterial genomes
 1662 and its consequences for bacterial community analyses. *PLoS ONE*, 8(2), e57923.
 1663 doi:10.1371/journal.pone.0057923
 1664 Visco, J. A., Apothélos-Perret-Gentil, L., Cordonier, A., Esling, P., Pillet, L., & Pawlowski, J. (2015).
 1665 Environmental Monitoring: Inferring the Diatom Index from Next-Generation Sequencing
 1666 Data. *Environmental Science and Technology*, 49(13), 7597–7605. doi:10.1021/es506158m
 1667 Vivien, R., Lejzerowicz, F., & Pawlowski, J. (2016). Next-generation sequencing of aquatic
 1668 oligochaetes: Comparison of experimental communities. *PLoS ONE*, 11(2), e0148644.
 1669 doi:10.1371/journal.pone.0148644
 1670 von Ammon, U., Wood, S. A., Laroche, O., Zaiko, A., Lavery, S. D., Inglis, G. J., & Pochon, X.
 1671 (2019). Linking environmental DNA and RNA for improved detection of the marine invasive
 1672 fanworm *Sabella spallanzanii*. *Frontiers in Marine Science*, 6, 621.

doi:10.3389/fmars.2019.00621

Washburne, A. D., Morton, J. T., Sanders, J., McDonald, D., Zhu, Q., Oliverio, A. M., & Knight, R. (2018). Methods for phylogenetic analysis of microbiome data. *Nature Microbiology*, 3(6), 652–661. doi:10.1038/s41564-018-0156-0

Waters, C. N., Zalasiewicz, J., Summerhayes, C., Barnosky, A. D., Poirier, C., Gałuszka, A., ... Wolfe, A. P. (2016). The Anthropocene is functionally and stratigraphically distinct from the Holocene. *Science*, 351(6269), aad2622. doi:10.1126/science.aad2622

Webb, C. O., Ackerly, D. D., McPeck, M. A., & Donoghue, M. J. (2002). Phylogenies and community ecology. *Annual Review of Ecology and Systematics*, 33(2002), 475–505. doi:10.1146/annurev.ecolsys.33.010802.150448

Weigand, H., Beermann, A. J., Čiampor, F., Costa, F. O., Csabai, Z., Duarte, S., ... Ekrem, T. (2019). DNA barcode reference libraries for the monitoring of aquatic biota in Europe: Gap-analysis and recommendations for future work. *Science of the Total Environment*, 678, 499–524. doi:10.1016/j.scitotenv.2019.04.247

Welsh, H. H., & Ollivier, L. M. (1998). Stream Amphibians as Indicators of Ecosystem Stress : A Case Study from California ' s Redwoods. *Ecological Applications*, 8(4), 1118–1132.

Wilcox, T. M., Carim, K. J., Young, M. K., McKelvey, K. S., Franklin, T. W., & Schwartz, M. K. (2018). Comment: The Importance of Sound Methodology in Environmental DNA Sampling. *North American Journal of Fisheries Management*, 38(3), 592–596. doi:10.1002/nafm.10055

Xu, Z., Malmer, D., Langille, M. G. I., Way, S. F., & Knight, R. (2014). Which is more important for classifying microbial communities: Who's there or what they can do? *ISME Journal*, 8(12), 2357–2359. doi:10.1038/ismej.2014.157

Yilmaz, P., Kottmann, R., Field, D., Knight, R., Cole, J. R., Amaral-Zettler, L., ... Glöckner, F. O. (2011). Minimum information about a marker gene sequence (MIMARKS) and minimum information about any (x) sequence (MlXS) specifications. *Nature Biotechnology*, 29(5), 415–420. doi:10.1038/nbt.1823

Zaiko, A., Samuiloviene, A., Ardura, A., & Garcia-Vazquez, E. (2015). Metabarcoding approach for nonindigenous species surveillance in marine coastal waters. *Marine Pollution Bulletin*,

1702 100(1), 53–59. doi:10.1016/j.marpolbul.2015.09.030

1703 Zappellini, C., Karimi, B., Foulon, J., Lacercat-Didier, L., Maillard, F., Valot, B., ... Chalot, M.

1704 (2015). Diversity and complexity of microbial communities from a chlor-alkali tailings

1705 dump. *Soil Biology and Biochemistry*, 90, 101–110. doi:10.1016/j.soilbio.2015.08.008

1706 Zhou, J., Deng, Y., Luo, F., Molecular, P., Network, E., Communities, S. M., ... Journal, A. S. M.

1707 (2011). Phylogenetic molecular ecological network of soil microbial. *MBio*, 2(4), 1–8.

1708 doi:10.1128/mBio.00122-11.Editor

1709 Zhou, X., Li, Y., Liu, S., Yang, Q., Su, X., Zhou, L., ... Huang, Q. (2013). Ultra-deep sequencing

1710 enables high-fidelity recovery of biodiversity for bulk arthropod samples without PCR

1711 amplification. *GigaScience*, 2(1), 4. doi:10.1186/2047-217X-2-4

1712 Zimmermann, J., Abarca, N., Enke, N., Enk, N., Skibbe, O., Kusber, W. H., & Jahn, R. (2014).

1713 Taxonomic reference libraries for environmental barcoding: a best practice example from

1714 diatom research. *PloS One*, 9(9), e108793. doi:10.1371/journal.pone.0108793

1715 Zinger, L., Bonin, A., Alsos, I. G., Bálint, M., Bik, H., Boyer, F., ... Taberlet, P. (2019). DNA

1716 metabarcoding—Need for robust experimental designs to draw sound ecological

1717 conclusions. *Molecular Ecology*, 28(8), 1857–1862. doi:10.1111/mec.15060

1718

1719